



PRIOR AUTHORIZATION POLICY

POLICY: Erectile Dysfunction – Sildenafil Prior Authorization Policy

- Viagra® (sildenafil tablets – Pfizer, generic)
- Vybriquetm (sildenafil oral film – IBSA Pharma)

REVIEW DATE: 11/12/2025; selected revision 02/11/2026, 03/04/2026

INSTRUCTIONS FOR USE

THE FOLLOWING COVERAGE POLICY APPLIES TO HEALTH BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. CERTAIN CIGNA COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. EACH COVERAGE REQUEST SHOULD BE REVIEWED ON ITS OWN MERITS. MEDICAL DIRECTORS ARE EXPECTED TO EXERCISE CLINICAL JUDGMENT WHERE APPROPRIATE AND HAVE DISCRETION IN MAKING INDIVIDUAL COVERAGE DETERMINATIONS. WHERE COVERAGE FOR CARE OR SERVICES DOES NOT DEPEND ON SPECIFIC CIRCUMSTANCES, REIMBURSEMENT WILL ONLY BE PROVIDED IF A REQUESTED SERVICE(S) IS SUBMITTED IN ACCORDANCE WITH THE RELEVANT CRITERIA OUTLINED IN THE APPLICABLE COVERAGE POLICY, INCLUDING COVERED DIAGNOSIS AND/OR PROCEDURE CODE(S). REIMBURSEMENT IS NOT ALLOWED FOR SERVICES WHEN BILLED FOR CONDITIONS OR DIAGNOSES THAT ARE NOT COVERED UNDER THIS COVERAGE POLICY (SEE "CODING INFORMATION" BELOW). WHEN BILLING, PROVIDERS MUST USE THE MOST APPROPRIATE CODES AS OF THE EFFECTIVE DATE OF THE SUBMISSION. CLAIMS SUBMITTED FOR SERVICES THAT ARE NOT ACCOMPANIED BY COVERED CODE(S) UNDER THE APPLICABLE COVERAGE POLICY WILL BE DENIED AS NOT COVERED. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

CIGNA NATIONAL FORMULARY COVERAGE:

OVERVIEW

Sildenafil (tablets [Viagra, generic] and oral film [Vybriquet]), a phosphodiesterase type 5 (PDE5) inhibitor, is indicated for the treatment of **erectile dysfunction**.^{1,17}

Sildenafil has been studied for other uses.

- **Benign Prostatic Hyperplasia.** The European Association of Urology guidelines (2024) note that phosphodiesterase type 5 inhibitors can be used in men with moderate-to-severe lower urinary tract symptoms with or without erectile dysfunction.⁹ The guidelines add that based on the results from a meta-analysis⁸, younger men with lower body mass index and more severe lower urinary tract symptoms benefit the most from phosphodiesterase type 5 inhibitors.
- **High-Altitude Pulmonary Edema (HAPE).** Published guidelines for the prevention and treatment of HAPE recommend nifedipine as the preferred pharmacologic treatment option.¹² Regarding PDE5 inhibitors, tadalafil is given a strong

recommendation for HAPE prevention in known susceptible individuals who are not candidates for nifedipine. Dexamethasone is recommended for prevention if the patient is not a candidate for nifedipine and tadalafil. For HAPE treatment, PDE5 inhibitors may be used if nifedipine is not available.

- **Prophylaxis after Radical Prostatectomy.** Sildenafil given on a daily basis has been used to improve the return of normal spontaneous erectile function, improve tissue oxygenation, and prevent penile fibrosis after nerve-sparing radical prostatectomy.^{10,11} It is better to initiate a penile rehabilitation program as soon as possible after surgery in order to limit and prevent postoperative local hypoxxygenation and fibrosis.
- **Pulmonary Arterial Hypertension (PAH).** PAH is a serious but rare condition affecting fewer than 20,000 patients in the US.^{13,14} It is classified within Group 1 pulmonary hypertension among the five different groups that are recognized by the World Health Organization (WHO). In this progressive disorder, the small arteries in the lungs become narrowed, restricted, or blocked causing the heart to work harder to pump blood, leading to activity impairment. Although the mean age of diagnosis is between 36 and 50 years, patients of any age may be affected, including pediatric patients. PAH is defined as a mean pulmonary artery pressure (mPAP) > 20 mmHg (at rest) with a pulmonary arterial wedge pressure (PAWP) ≤ 15 mmHg and a pulmonary vascular resistance > 2 Wood units measured by cardiac catheterization.¹⁵

The CHEST guideline and Expert Panel Report regarding therapy for PAH in adults (2019) details many medications. It was noted that PDE5 inhibitors play a vital role and have various benefits in the management of PAH.¹⁴ The European Society of Cardiology and the European Respiratory Society guidelines regarding the treatment of pulmonary hypertension (2022) also recognize PDE5 inhibitors as having a prominent role in the management of this condition, as monotherapy or in use as combination with other agents.¹⁶

Sildenafil tablets, suspension, and injection (Revatio[®], generics) are approved for pulmonary arterial hypertension.² Sildenafil (Viagra, generics) are available in 25 mg, 50 mg, and 100 mg tablets, and Revatio is available as 20 mg tablets. Viagra has been used for this diagnosis.^{3,4} Doses of Viagra that were used in these reports ranged from 25 mg twice daily to 100 mg five times daily. Patients were typically started on Revatio 20 mg three times daily.

- **Raynaud's Phenomenon.** There are studies which show sildenafil has been beneficial in patients with Raynaud's phenomenon.^{5,6} Guidelines from the European League against Rheumatism (EULAR) on the treatment of systemic sclerosis (2023) recommend considering dihydropyridine calcium channel blockers (CCBs), usually oral nifedipine, for first-line therapy of Raynaud's phenomenon in patients with systemic sclerosis.⁷ Phosphodiesterase type 5 inhibitors should also be considered in such clinical scenarios.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of sildenafil. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with sildenafil as well as the monitoring required for adverse events and long-term efficacy, some approvals require sildenafil to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Documentation: Documentation is required for initiation of therapy as noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart

notes and catheterization laboratory reports. All documentation must include patient-specific identifying information. For a patient case in which the documentation requirement of the right heart catheterization upon Prior Authorization coverage review for a different medication indicated for pulmonary arterial hypertension (PAH) World Health Organization (WHO) Group 1 has been previously provided, the documentation requirement in this *Erectile Dysfunction – Sildenafil Prior Authorization Policy* is considered to be met.

- **Viagra® (sildenafil tablets - Pfizer, generic)**
- **Vybrique™ (sildenafil oral film – IBSA Pharma)**

is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indication

1. Erectile Dysfunction. Approve for 1 year.

Other Uses with Supportive Evidence

2. Benign Prostatic Hyperplasia. Approve for 1 year if the patient meets ONE of the following (A or B):

Note: For men with erectile dysfunction and benign prostatic hyperplasia, use criterion 1 above.

A) Patient has tried an alpha-1 (α 1) blocker; OR

Note: Examples of alpha-1 (α 1) blockers include doxazosin, terazosin, tamsulosin, alfuzosin.

B) Patient has tried a 5 α -reductase inhibitor.

Note: Examples of a 5 α -reductase inhibitor include finasteride and dutasteride.

3. High-Altitude Pulmonary Edema (HAPE), Treatment or Prevention. Approve for 1 year if the patient meets BOTH of the following (A and B):

A) Patient has HAPE or a history of HAPE; AND

B) Patient has tried one other pharmacologic therapy for the treatment or prevention of HAPE.

Note: Examples of other pharmacologic therapy for the treatment or prevention of HAPE are nifedipine, dexamethasone, and Cialis (tadalafil tablets).

4. Prophylaxis After Radical Prostatectomy (Early Penile Rehabilitation). Approve for 1 year if the patient meets BOTH of the following (A and B):

A) Patient had radical prostatectomy within the previous 12 months; AND

B) The medication is prescribed by or in consultation with a urologist.

5. Pulmonary Arterial Hypertension (PAH) [World Health Organization {WHO} Group 1]. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 1 year if the patient meets BOTH of the following (i and ii):

i. Patient meets BOTH of the following (a and b):

a) Patient has had a right heart catheterization **[documentation required]**; AND

b) Results of the right heart catheterization confirm the diagnosis of WHO Group 1 PAH; AND

ii. The medication is prescribed by or in consultation with a cardiologist or a pulmonologist; OR

- B) Patient is Currently Receiving Sildenafil for PAH.** Approve for 1 year if the patient meets BOTH of the following (i and ii):
- i. Patient meets BOTH of the following (a and b):
 - a) Patient has had a right heart catheterization; AND
Note: This refers to prior to starting therapy with a medication for WHO Group 1 PAH.
 - b) Results of the right heart catheterization confirm the diagnosis of WHO Group 1 PAH; AND
 - ii. The medication is prescribed by or in consultation with a cardiologist or a pulmonologist.

6. Raynaud's Phenomenon. Approve for 1 year if the patient meets ONE of the following (A or B):

- A)** Patient has tried one calcium channel blocker; OR
Note: Examples of calcium channel blockers include amlodipine, felodipine, nifedipine.
- B)** According to the prescriber, use of a calcium channel blocker is contraindicated.
Note: Examples of reasons a patient cannot take calcium channel blocker therapy include right heart failure or decreased cardiac output.

CONDITIONS NOT COVERED

- **Viagra® (sildenafil tablets - Pfizer, generic)**
- **Vybriq™ (sildenafil oral film - IBSA Pharma)**

is(are) considered not medically necessary for ANY other use(s) ; criteria will be updated as new published data are available).

REFERENCES

1. Viagra® tablets [prescribing information]. New York, NY: Pfizer; December 2017.
2. Revatio® tablets [prescribing information]. New York, NY: Pfizer; January 2023.
3. Garg N, Sharma MK, Sinha N. Role of oral sildenafil in severe pulmonary arterial hypertension: Clinical efficacy and dose response relationship. *Int J Cardiol.* 2007;120:306-313.
4. Galié N, Ghofrani HA, Torbicki A, et al; Sildenafil Use in Pulmonary Arterial Hypertension (SUPER) Study Group. Sildenafil citrate therapy for pulmonary arterial hypertension. *N Engl J Med.* 2005;353:2148-2157.
5. Fernandez-Codina A, Canas-Ruano E, Pope JE. Management of Raynaud's phenomenon in systemic sclerosis-a practical approach. *J Scleroderm Relat Disord.* 2019;4(2):102-110.
6. Hinze AM, Wigley FM. Pharmacotherapy options in the management of Raynaud's phenomenon. *Curr Treat Opt Rheumatol.* 2018;4(3):235-254.
7. Del Galdo F, Lescoat A, Conaghan PG, et al. 2023 Update of EULAR recommendations for the treatment of systemic sclerosis. *Ann Rheum Dis.* 2023;82:154-155.
8. Gacci M, Corona G, Salvi M, et al. A systematic review and meta-analysis on the use of phosphodiesterase 5 inhibitors alone or in combination with α -blockers for lower urinary tract symptoms due to benign prostatic hyperplasia. *Eur. Urol.* 2012;61:994-1003.
9. Cornu JH, Gacci M, Hashim H, et al. Guidelines on the management of non-neurogenic male lower urinary tract symptoms (LUTS). © European Association of Urology 2024. Available at: <https://uroweb.org/guidelines/management-of-non-neurogenic-male-luts>. Accessed on November 6, 2025.
10. Bannowsky A, Schulze H, van der Horst C, et al. Recovery of erectile function after nerve-sparing radical prostatectomy: improvement with nightly low-dose sildenafil. *BJU Int.* 2008;101:1279-1283.

11. Bratu O, Oprea I, Marcu D, et al. Erectile dysfunction post-radical prostatectomy- a challenge for both patient and physician. *J Med Life*. 2017;10(1):13-18.
12. Luks AM, Beidleman BA, Freer L, et al. Wilderness Medical Society Clinical Practice Guidelines for the Prevention, Diagnosis, and Treatment of Acute Altitude Illness: 2024 Update. *Wilderness Environ Med*. 2024 Mar;35(1):S2-S19.
13. Ruopp NF, Cockrill BA. Diagnosis and treatment of pulmonary arterial hypertension. A review. *JAMA*. 2022;327(14):1379-1391.
14. Klinger JR, Elliott CG, Levine DJ, et al. Therapy for pulmonary arterial hypertension in adults. Update of the CHEST guideline and Expert Panel Report. *CHEST*. 2019;155(3):565-586.
15. Maron BA. Revised Definition of Pulmonary Hypertension and Approach to Management: A Clinical Primer. *J Am Heart Assoc*. 2023 Apr 18;12(8):e029024. [Epub].
16. Humbert M, Kovacs G, Hoeper MM, et al, for the ESC/ERS Scientific Document Group. 2022 ESC/ERS guidelines for the diagnosis and treatment of pulmonary hypertension. *Eur Heart J*. 2022;43(38):3618-3731.
17. Vybrique™ oral film [prescribing information]. Parsippany, NJ: IBSA Pharma; December 2025.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	No criteria changes.	11/01/2023
Annual Revision	No criteria changes.	11/06/2024
Annual Revision	No criteria changes.	11/12/2025
Selected Revision	High-Altitude Pulmonary Edema (HAPE), Treatment or Prevention: Serevent (salmeterol inhalation powder) and acetazolamide were removed as examples of pharmacologic therapy for the treatment or prevention of HAPE. Pulmonary Arterial Hypertension (PAH): Added that PAH is World Health Organization (WHO) Group 1. For Initial Therapy and a Patient Currently Receiving Sildenafil for PAH, requirements were added that the patient has had a right heart catheterization (documentation required for Initial Therapy) and the results confirm the diagnosis of WHO Group 1 PAH, and that the medication is prescribed by or in consultation with a cardiologist or a pulmonologist.	02/11/2026
Selected Revision	Vybrique (sildenafil oral film) was added to the policy with the same authorization criteria and exclusions as other sildenafil products.	03/04/2026

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2026 The Cigna Group.