



Medical Coverage Policy

Effective Date4/15/2026

Next Review Date4/15/2027

Coverage Policy Number..... 0309

Surgical Treatment of Chest Wall Deformities

Table of Contents

- Overview 2
- Coverage Policy 2
- Coding Information 3
- General Background..... 4
- Health Equity Considerations..... 10
- References..... 10
- Revision Details 14

Related Coverage Resources

- [Bone Graft Substitutes](#)
- [Breast Reconstruction Following Mastectomy or Lumpectomy](#)
- [Breast Reduction](#)
- [Chest Imaging Guidelines](#)
- [Transthoracic Echocardiography in Adults](#)
- [Transthoracic Echocardiography in Children](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer’s particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer’s benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer’s benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted

for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses surgical repair of chest wall deformities in pectus excavatum, pectus carinatum, and Poland syndrome.

Coverage Policy

Coverage for surgical repair of a chest wall deformity varies across plans. Refer to the customer's benefit plan document for coverage details.

Pectus Excavatum

Surgical repair of pectus excavatum is considered medically necessary when imaging studies (e.g., computerized tomography [CT] scans, radiographs, magnetic resonance imaging [MRI]) confirm a pectus index (i.e., Haller index) greater than 3.25 and EITHER of the following criteria is met:

- pulmonary function studies demonstrate restrictive or obstructive lung disease
- cardiac imaging (e.g., echocardiography, stress echocardiography, MRI) demonstrates findings consistent with external cardiac compression

Magnetic Mini Mover Procedure (3MP) for the treatment of pectus excavatum is considered experimental, investigational or unproven.

Pectus Carinatum

Surgical repair of pectus carinatum is considered medically necessary when there is documented evidence of significant physical functional impairment (e.g., cardiac or respiratory insufficiency), and the procedure is expected to correct the impairment.

Poland Syndrome

Surgical treatment of a chest wall deformity associated with Poland syndrome, including initial breast reconstruction*, is considered medically necessary when EITHER of the following criteria is met:

- preoperative frontal photographs demonstrate partial or complete absence of the pectoralis major muscle(s)
- preoperative imaging demonstrates absence or hypoplasia of upper costal cartilage and/or ribs

***Note:** Please reference the Cigna Medical Coverage policies "Breast Reconstruction Following Mastectomy or Lumpectomy" and "Breast Reduction" for conditions of coverage related specifically to breast reconstruction procedures.

When the above criterion is met, coverage of initial breast reconstruction of the affected breast is limited to a one- or two-stage approach (e.g., direct-to-implant or tissue expander followed by permanent implant) to correct breast asymmetry.

Breast reconstruction of the contralateral breast to achieve symmetry in the absence of a severe physical deformity or disfigurement of the contralateral breast accompanied by functional deficit is considered cosmetic in nature and not medically necessary.

Breast reconstruction revision surgery of the affected breast following initial treatment of Poland syndrome is considered medically necessary when there is severe physical deformity or disfigurement accompanied by functional deficit.

Breast reconstruction revision surgery following initial treatment of Poland syndrome that is performed for the sole purpose of improving appearance is considered cosmetic in nature and not medically necessary.

When performed for indications other than those listed above, surgical repair of a chest wall deformity is considered not medically necessary.

Coding Information

Notes:

1. This list of codes may not be all-inclusive since the American Medical Association (AMA) and Centers for Medicare & Medicaid Services (CMS) code updates may occur more frequently than policy updates.
2. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Pectus Carinatum/Pectus Excavatum

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
21740	Reconstructive repair of pectus excavatum or carinatum; open
21742	Reconstructive repair of pectus excavatum or carinatum; minimally invasive approach (Nuss procedure), without thoracoscopy
21743 [†]	Reconstructive repair of pectus excavatum or carinatum; minimally invasive approach (Nuss procedure), with thoracoscopy

[†]Note: Considered Experimental/Investigational/Unproven when used to report Magnetic Mini Mover (3MP) procedure.

Poland Syndrome

Considered Medically Necessary when used to report surgical repair of a chest deformity associated with Poland syndrome:

CPT®* Codes	Description
15771	Grafting of autologous fat harvested by liposuction technique to trunk, breasts, scalp, arms, and/or legs; 50 cc or less injectate
15772	Grafting of autologous fat harvested by liposuction technique to trunk, breasts, scalp, arms, and/or legs; each additional 50 cc injectate, or part thereof (List separately in addition to code for primary procedure)
15777	Implantation of biologic implant (eg, acellular dermal matrix) for soft tissue reinforcement (ie, breast, trunk) (List separately in addition to code for primary procedure)
19325	Breast augmentation with implant
19357	Tissue expander placement in breast reconstruction, including subsequent expansion(s)
19380	Revision of reconstructed breast (eg, significant removal of tissue, re-advancement and/or re-inset of flaps in autologous reconstruction or significant capsular revision combined with soft tissue excision in implant-based reconstruction)

***Current Procedural Terminology (CPT®) ©2025 American Medical Association: Chicago, IL.**

General Background

The thorax (i.e., chest cavity) is a rigid structure that protects the thoracic organs and supports the upper extremities. Commonly reported chest wall deformities include pectus excavatum (PE), pectus carinatum (PC), and Poland syndrome. In many cases, primarily cosmetic complaints without functional impairment are associated with the abnormality. However, abnormalities of the chest wall can lead to restrictive or obstructive pulmonary disease, impaired respiratory muscle strength, and decreased ventilatory performance in response to physical stress (Boas and Kier, 2025). Cardiac or respiratory impairment may result in functional limitations, such as activity intolerance. Other symptoms may include mild to moderate exercise limitation, respiratory infections, and asthmatic symptoms, as well as decreased stamina and endurance.

Pectus Excavatum

Pectus excavatum (PE), also referred to as a sunken chest or funnel chest, is the most common congenital chest wall deformity. The deformity may be deeper on the right side than on the left side and result in a rotation of the sternum. It is usually diagnosed within the first year of life, with wide variations in the degree of sternal depression. PE is three to four times more prevalent in males, and occurs most frequently in Caucasians (Mourisse and van der Heide, 2022; Gottlieb, et al., 2018). Although most patients are asymptomatic, during periods of rapid bone growth (e.g., puberty), the appearance of the chest may worsen and symptoms may develop. Moderate to severe deformities may displace the heart into the left chest, decreasing stroke volume and cardiac output. A 2024 systematic review and meta-analysis found that individuals with PE were more than six times more likely to have mitral valve prolapse, compared to individuals without PE (Sonaglioni, et al., 2024). Chest deformities may also depress the sternal volume, adversely affecting the flow of air in and out of the lungs. Scoliosis, congenital heart disease, and functional heart murmurs can also be associated with PE.

Symptoms of PE may include fatigue, dyspnea, chest discomfort, and palpitations with mild exercise. The body generally compensates by increasing the heart rate with activity to overcome the decreased cardiac output and by more rapid, shallow breathing to compensate for the respiratory deficit. Pulmonary effects associated with PE generally include restrictive lung disease,

atelectasis, and paradoxical respiration. Restricted lung disease is evident when the total lung capacity (TLC) (forced vital capacity plus residual capacity) is less than 80% of the predicted value for an individual (Boas and Kier, 2025; Mourisse and van der Heide, 2022; Rust and Westney; 2016).

Pectus Carinatum

Pectus carinatum (PC) (i.e., pigeon breast or chicken breast) is a congenital chest deformity characterized by an anterior protrusion deformity of the sternum and costal cartilages. The condition is more rare than PE, and is six times more prevalent in males than in females (Gottlieb, et al.; 2018). PC is typically not confirmed until after the growth spurts of early adolescence. This deformity produces a rigid chest and, while symptoms are uncommon, it may result in inefficient respiration as a result of the restrictive chest formation. Three types of PC-related defects have been identified in the literature:

- anterior displacement of the body of the sternum and symmetrical concavity of the costal cartilages
- lateral depression of the ribs on one or both sides of the sternum
- pouter pigeon breast (the least common of the three): a defect that consists of an upper or chondromalacial prominence with protrusion of the manubrium and depression of the sternal body

The degree of physiological impairment is related to the degree of chest deformity. Patients with PC may develop symptoms as a result of restricted air exchange; complete expiration of air from the lungs may not occur. In addition, pain may result from the secondary pressures that develop from the overgrowth of cartilage. Other conditions that may be associated with PC include frequent respiratory infections, asthma, rickets, Marfan's syndrome, scoliosis, mitral valve disease, and other cardiac changes.

Non-surgical treatment may include orthotic bracing. Bracing may be effective for correction of the chest wall deformity associated with PC, however this type of treatment is often aimed at improving cosmesis. Compression orthotic braces exert pressure on the anteroposterior direction and may be recommended for skeletally immature children (e.g., < age 18 years) with mild to moderate chest deformities. Prolonged use of the orthotic device is often required for total correction, and compliance is an important factor for successful remodeling. As a result candidates must be motivated to wear the brace (Gottlieb, et al., 2018).

Poland Syndrome

Poland syndrome (i.e., Poland's anomaly, Poland's syndactyly), is a rare congenital disorder characterized by the absence or hypoplasia of the sternocostal head of the pectoralis major muscle, associated with a wide range of other malformations. Co-occurring anomalies include absence or hypoplasia of the pectoralis minor muscle, absence or underdevelopment the upper ribs, absence of costal cartilages, hypoplasia of the breast and subcutaneous tissue, and a variety of hand and upper-extremity anomalies. The right side of the body is affected twice as often as the left, and the condition occurs more frequently in men than in women. When the anomaly occurs on the left side of the body, the heart and lungs are vulnerable, because they may be covered only by skin, fascia and pleura. In cases of severe cartilage deficiency, patients may develop lung hernia and paradoxical respiratory motion. In less severe cases, patients may develop a simple flattening of the anterior chest wall.

Diagnosis and Evaluation

The severity of the chest wall abnormality is dependent upon the depth, symmetry and width of the deformity. Chest radiographs are commonly used to determine the degree of chest wall deformity. Cross-sectional imaging such as computed tomography (CT) scans and magnetic

resonance imaging (MRI) may be used to evaluate the degree of cardiac compression (ventricular compression), pulmonary compression, and cardiac displacement. Plain anteroposterior and lateral radiographs, CT and MRIs are used to determine the Haller index (a measurement of chest diameter). Scan ratios that reveal transverse to AP diameter of greater than 3.25 are considered significant for pectus excavatum. A normal chest has an index of 2.5 (Mourisse and van der Heide, 2022). The Correction Index (CI) is another, more recent thoracic index used to assess the severity of deformity in pectus excavatum. The CI corresponds to the depth of the anterior wall depression, reflected as a percentage of the sternal/costal cartilage depression to be corrected with surgery. There is no widely accepted CI cut-off point for surgical repair, though some authors have suggested surgical correction may be appropriate for symptomatic individuals with a CI \geq 28% (Zens, et al., 2022; Rodriguez-Granillo, et al., 2019; Sujka, et al., 2018; Poston, et al., 2014; St. Peter, et al., 2011). The Haller index remains the gold standard for estimating the degree of chest wall deformity in pectus excavatum.

Echocardiography and/or electrocardiography may also be used to evaluate cardiac status. Respiratory status can be determined with the use of pulmonary function studies. In some cases, pulmonary function studies may reveal a restrictive pattern (incomplete lung expansion) and a subsequent decrease in pulmonary volume and reserve. The forced expiratory volume (in one second) (FEV₁), forced vital capacity (FVC), and total lung capacity (TLC) are reduced while the ratio of FEV₁/FVC may be normal or increased in the presence of restrictive airway disease. TLC <80% predicted value signifies restrictive pulmonary disease (Rust and Westney; 2016).

The diagnosis of Poland syndrome is usually obtained by clinical examination. Chest wall abnormalities and determining the presence of latissimus dorsi muscles may require CT scans while chest radiographs may be utilized to evaluate rib formation.

Surgical Treatment

Indications for surgical correction vary widely. Surgical repair is offered primarily as a method of improving cosmesis and psychological factors but may be necessary to improve cardiopulmonary function in some patients, as the disfigurement may be accompanied by physiologic impairment.

Pectus Excavatum/Pectus Carinatum: While the optimal age for surgical repair is generally between the ages of 11 and 18 years, and may be performed in adults, each case must be reviewed individually for the presence of cardiopulmonary impairment. Criteria that may be used to demonstrate severe PE and the need for surgical repair include two or more of the following (Mourisse and van der Heide, 2022):

- a Haller CT index greater than 3.25
- pulmonary function studies that indicate restrictive or obstructive airway disease
- a cardiology evaluation, where the chest compression is causing murmurs, mitral valve prolapse, cardiac displacement, or conduction abnormalities on the echocardiogram or electrocardiogram (ECG)
- documentation of progression of the deformity with associated physical symptoms other than isolated concerns of body image
- a failed Ravitch procedure
- a failed minimally invasive procedure

Surgery for PE may be performed using any one of several techniques, including a sternal osteotomy (i.e., a modified osteotomy that involves supporting, removing and repositioning the sternum) or implantation of a Silastic mold in the subcutaneous space to fill the defect without altering the thoracic cage. Surgical correction often employs a metal bar behind the sternum; the bar may be removed in one to two years, after remodeling has occurred. The standard surgical procedure is the open Ravitch procedure, which involves extensive dissection, cartilage resection

and sternal osteotomy. More recently, minimally invasive techniques, such as the Nuss procedure (i.e., a minimally invasive repair of pectus excavatum [MIRPE]), have been utilized. These procedures involve the insertion of a convex steel bar beneath the sternum through small thoracic incisions and do not require cartilage resection or osteotomy. It has been reported that the Nuss procedure has a shorter operative time and less operative blood loss compared to the Ravitch method (Mao, et al., 2017).

Surgical repair may also be recommended for the treatment of PC. Surgical repair of PC may involve removing the affected cartilages and mobilizing the skin and pectoralis muscle flaps. To straighten the sternum, any one of the following surgeries may be performed:

- an osteotomy
- a subperichondrial resection of the involved costal cartilages
- a wedge-shaped osteotomy in the anterior sternal plate

Literature Review

Published evidence evaluating surgical repair of chest wall deformities consists of meta-analyses, case series, cross-comparison studies, and prospective trials. The reported clinical outcomes are mixed; differences among outcomes may be related to patient selection criteria, the degree of severity of the deformity, the surgical technique utilized, and growth effects. Many authors evaluated and reported on the methods of surgical repair and improved cosmetic outcomes while some evaluated the impact of PE or PC on cardiopulmonary function. There is little consensus regarding the degree of cardiopulmonary impairment, if any, that is associated with these anomalies. The effects of surgery on exercise tolerance are not clearly established, however, data suggesting improvement in cardiovascular and/or pulmonary function and activity tolerance after surgical repair have been reported (Sonaglioni, et al., 2023; Kelly, et al., 2013; Jaroszewski and Fonkalsrud, 2007; Kubiak, et al., 2007; Lawson, et al., 2005; Bawazir, et al., 2005; Fonkalsrud and Anselmo, 2004; Haller and Loughlin, 2000; Fonkalsrud, et al., 1994). Outcome measures of these studies generally included total lung capacity (TLC), functional residual capacity (FRC), vital capacity (VC), expiratory flow rate (EFR), and maximum expiratory flow rate (MEFR), exercise tolerance and endurance typically measured prior to surgery, immediately following surgery and three to six months postoperatively. Improvement was generally seen only with increased periods of exercise and not with routine pulmonary function testing at rest. The results of some meta-analyses and other published clinical studies in the medical literature are also mixed, some results tend to support improvement in cardiopulmonary function following surgery (Johnson, et al., 2008; Malek, et al., 2006a, Malek, et al., 2006b) while others do not (Zganjer and Zganjer, 2010; Castellani, et al., 2010; Guntheroth and Spiers, 2007).

Surgical repair of chest wall deformities, such as pectus excavatum and pectus carinatum, is widely recognized as an accepted and established treatment option within contemporary clinical practice. The Nuss procedure for pectus excavatum and the anti-Nuss procedure or Abramson technique for pectus carinatum are described in major surgical and medical textbooks as operative approaches for appropriately selected patients. Their inclusion in these references reflects broad consensus among thoracic surgeons, pediatric surgeons, and multidisciplinary specialists that minimally invasive bar-based repair techniques are effective, well-established corrective options for these congenital chest wall deformities (Fillmore, 2026; Lovegreen and Mogensen, 2026; Graham and Sanchez-Lara, 2025; Demehri, 2024).

Poland Syndrome: Patients with Poland syndrome typically present for surgical reconstruction to correct deformity or defect of the pectoralis muscles and/or ribs, and to correct breast asymmetry. In cases of rib aplasia, operative reconstruction may eliminate paradoxical motion, improving respiratory impairment. For more severe conditions, reconstructive surgery also provides protection of the underlying heart and lung structures. Surgical repair of the chest wall includes

the reconstruction of the pectoral muscles and resection of deformed cartilages. This repair typically involves muscle transfers and/or flaps to match normal development of the unaffected side, reconstruction of the axillary line, and correction of infraclavicular flattening. Techniques for repair of large defects include the use of rib grafts, other bony allo- or autografts, mesh, or a combination of these techniques (Gottlieb, et al., 2018; Fokin and Robicsek; 2002).

Surgical treatment of Poland syndrome may include reconstruction of the breast and nipple on the affected side by a plastic surgeon, in addition to surgical repair of the chest wall muscles and hypoplastic bone. Surgery may be performed early (approximately age 13) in males, however, in females, reconstructive surgery is often deferred until breast development is complete. If there are rib abnormalities and paradoxical motion, the rib grafts or other chest wall stabilization may occur before breast development is complete. Generally, reconstruction of the breast involves tissue expansion, placement of permanent breast implants, and may involve myocutaneous or latissimus dorsi flaps if there is an associated anomaly of the pectoral muscle. Nipple-areolar reconstruction is generally performed at a later stage. Consequently, for patients with Poland syndrome, treatment provided before complete breast development may involve the use of tissue expanders in the affected side which can be inflated periodically to match development of the unaffected breast. Expanders allow for tissue expansion and accommodation of a permanent implant and latissimus muscle upon completion of breast development. Once breast development is complete, the expander is removed, a permanent prosthesis is inserted, and breast reconstruction is performed.

For information on breast prostheses and products used in breast reconstructive procedures, please see Cigna Medical Coverage policy "Breast Reconstruction Following Mastectomy or Lumpectomy".

U.S. Food and Drug Administration (FDA)

Reconstructive repair of chest wall deformities may involve the placement of metal bars, struts, and stabilizers to correct the defect. These appliances are categorized as Class II medical devices, are regulated by the FDA via the 510(k) pathway, and are indicated as bone fixation devices for repair of PE and other anterior chest wall deformities.

Device or Product	Identifier	Manufacturer
KLS Martin Pure Pectus System	K250988	KLS-Martin L.P.
Lorenz Pectus Support Bar System	K061384	Walter Lorenz Surgical, Inc.
Park’s Pectus System	K191057	TDM Co. Ltd.
Pectus Versa System	K251050	GM Doz Reis Industria e Comercio

*FDA product codes: HRS

Note: Coverage decisions are not based solely on FDA approval. Device or product names are provided for example purposes only. Their inclusion does not indicate endorsement or preference for any specific brand or model. This list is not intended to reflect all available products or technologies.

Literature Review

Pectus Excavatum and Pectus Carinatum: Published evidence evaluating surgical repair of chest wall deformities consists of meta-analyses, case series, cross-comparison studies, and prospective trials. The reported clinical outcomes are mixed; differences among outcomes may be related to patient selection criteria, the degree of severity of the deformity, the surgical technique

utilized, and growth effects. Many authors evaluated and reported on the methods of surgical repair and improved cosmetic outcomes while some evaluated the impact of PE or PC on cardiopulmonary function. There is little consensus regarding the degree of cardiopulmonary impairment, if any, that is associated with these anomalies. The effects of surgery on exercise tolerance are not clearly established, however, data suggesting improvement in cardiovascular and/or pulmonary function and activity tolerance after surgical repair have been reported (Sonaglioni, et al., 2023; Kelly, et al., 2013; Jaroszewski and Fonkalsrud, 2007; Kubiak, et al., 2007; Lawson, et al., 2005; Bawazir, et al., 2005; Fonkalsrud and Anselmo, 2004; Haller and Loughlin, 2000; Fonkalsrud, et al., 1994). Outcome measures of these studies generally included total lung capacity (TLC), functional residual capacity (FRC), vital capacity,(VC), expiratory flow rate (EFR), and maximum expiratory flow rate (MEFR), exercise tolerance and endurance typically measured prior to surgery, immediately following surgery and three to six months postoperatively. Improvement was generally seen only with increased periods of exercise and not with routine pulmonary function testing at rest. The results of some meta-analyses and other published clinical studies in the medical literature are also mixed, some results tend to support improvement in cardiopulmonary function following surgery (Johnson, et al., 2008; Malek, et al., 2006a, Malek, et al., 2006b) while others do not (Zganjer and Zganjer, 2010; Castellani, et al, 2010; Guntheroth and Spiers, 2007).

Surgical repair of chest wall deformities, such as pectus excavatum and pectus carinatum, is widely recognized as an accepted and established treatment option within contemporary clinical practice. The Nuss procedure for pectus excavatum and the anti-Nuss procedure or Abramson technique for pectus carinatum are described in major surgical and medical textbooks as operative approaches for appropriately selected patients. Their inclusion in these references reflects broad consensus among thoracic surgeons, pediatric surgeons, and multidisciplinary specialists that minimally invasive bar-based repair techniques are effective, well-established corrective options for these congenital chest wall deformities (Fillmore, 2026; Lovegreen and Mogensen, 2026; Graham and Sanchez-Lara, 2025; Demehri, 2024).

Magnetic Mini-Mover Procedure (3MP): A procedure that is currently under investigation for the treatment of PE is the Magnetic Mini-Mover Procedure (3MP) which utilizes a magnetically coupled implant to pull the sternum forward and remodel the deformed costal cartilage. The internal magnetic implant is surgically placed on the sternum and an external magnet is applied using a custom-fitted chest wall orthotic device. The Magnetic Mini-Mover device was the subject of a United States Food and Drug Administration (FDA)-sponsored Phase III Investigational Device Exemption (IDE) clinical trial. IDE approval allows the device to be used in a clinical study for the collection of safety and effectiveness data. Current evidence evaluating the magnetic mini mover procedure in the published peer-reviewed scientific literature is primarily in the form of pilot studies and small case series (n=15). Reported adverse events have included postoperative pneumothorax, surgical site infection, and device breakage occurred in approximately half of the cases (Graves, et al., 2017; Harrison, et al., 2012; Harrison, et al., 2007).

Poland Syndrome: Surgical treatment for chest wall deformities associated with Poland syndrome is considered medically necessary because the condition involves congenital absence, underdevelopment, or agenesis of chest wall structures. Preoperative photographs are typically sufficient to establish the diagnosis based on characteristic anatomic asymmetry and absence of musculature. Once the condition has been diagnosed, surgical correction of the resulting breast asymmetry is limited to a one- or two-stage reconstructive approach (e.g., direct-to-implant placement or tissue expansion followed by a permanent implant). This reconstructive pathway is also supported in surgical and medical textbooks as treatment approaches for appropriately selected individuals. Inclusion in these references reflects broad consensus among plastic and pediatric surgeons. Surgical reconstruction in this context is performed to restore normal anatomical form and function, address asymmetry, and support physical and psychosocial

well-being, aligning with the definition of medically necessary reconstructive surgery (Beh, et al., 2024; Khan, et al., 2024; Cugno and Azzi, 2022; Muthialu, et al., 2022).

Professional Societies/Organizations

American Pediatric Surgical Association (APSA): In 2012, the APSA published its guideline on the evaluation and management of pectus carinatum. The guideline recommendations were based upon a systematic review of medical literature and peer-reviewed journals in or after 1973. Recommendations pertaining to surgical intervention included:

- Operative therapy plans should be individualized and age-appropriate, with surgery typically being deferred until pubertal growth is nearly complete.
- Open surgical techniques are acceptable for repair when performed by experienced pediatric surgeons.
- Thoracoscopic and other minimally invasive techniques are acceptable in some children, when performed by pediatric surgeons with advanced skills and experience in the technique.
- Surgery in children ages 5 years and younger is discouraged due to the risk of disruption of normal chest wall growth and chest wall restriction.
- The pediatric surgeon should be involved in extended postoperative follow up.

Health Equity Considerations

Health equity is the highest level of health for all people; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which people are born, grow, live, work, and age.

Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include safe housing, transportation, and neighborhoods; racism, discrimination and violence; education, job opportunities and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

Pitt et al. (2023) conducted an observational analysis of a tertiary pediatric multispecialty chest wall deformity clinic to describe the demographics of children aged 0–18 years who presented for initial evaluation between 2017 and 2021 and were diagnosed according to International Classification of Diseases criteria. Variables included diagnosis, age at first visit, sex, race/ethnicity, whether surgical correction of pectus excavatum (SCOPE) occurred during the study period, and patient-reported characteristics (worsening deformity, shortness of breath, palpitations, sports participation, and decreased physical activity). Within the 1510-child cohort, 50.0% (n=755) had pectus excavatum and 43.3% (n=653) had pectus carinatum; Poland syndrome accounted for 0.7% (n=10). Among pectus excavatum patients, 239 underwent SCOPE (31.7%). The operation rate was significantly higher in males than females (34.3% versus 22.3%; $P=0.003$), did not differ significantly by race/ethnicity ($p=0.158$), and age at SCOPE did not significantly differ by sex ($p=0.716$). Patient-reported worsening deformity, decreased physical activity, shortness of breath, and palpitations were each associated with higher likelihood of SCOPE ($p < 0.05$ for all listed symptoms). No Poland syndrome patients underwent surgical intervention; one was treated with bracing. The authors note single-institution and regional-demographics limitations and conclude that demographic distributions may differ from historical reports, while standardized evaluation in a dedicated clinic may mitigate differences in treatment (Pitt, et al., 2023).

References

1. American Pediatric Surgical Association (APSA). Pectus Carinatum Guideline. 2012. Accessed Jan 30, 2026. Available at URL address: <https://apsaped surg.org/resources/resources/practice-guidelines/apsa-guidelines/>
2. Bawazir OA, Montgomery M, Harder J, Sigalet DL. Midterm evaluation of cardiopulmonary effects of closed repair of pectus excavatum. *J Pediatr Surg.* 2005 May;40(5):863-7.
3. Beh HZ, Ferry AM, Dibbs RP, Buchanan EP, Monson LA. Pediatric chest and trunk deformities. In: Losee JE, Hopper RA. *Plastic Surgery, volume 3: Craniofacial, head, and neck surgery and pediatric plastic surgery.* Philadelphia, PA: 2024. 32, 974-987.
4. Boas S, Kier C. Skeletal Diseases Influencing Pulmonary Function. In: Kliegman RM, St Geme JW, Blum NJ, Tasker RC, Wilson KM, Schuh AM, Mack CL, editors; *Nelson Textbook of Pediatrics, 22nd ed.* 2025. Chapter 467, 2713-27115.e2
5. Castellani C, Windhaber J, Schober PH, Hoellwarth ME. Exercise performance testing in patients with pectus excavatum before and after Nuss procedure. *Pediatr Surg Int.* 2010 Jul;26(7):659-63.
6. Cugno S, Azzi AJ. Congenital breast anomalies. In: Farhadieh RD, Bulstrode NW, Mehrara BJ, Cugno S. *Plastic surgery: principles and practice.* Philadelphia, PA: Elsevier Inc. 2022. 37, 565-573.
7. Demehri FR and Shamberger RC. Congenital chest wall deformities. In: Sellke FW, del Nido PJ, Swanson SJ. *Sabiston and Spencer surgery of the chest.* 10th ed. Philadelphia, PA: Elsevier; 2024. 24, 453-486.
8. Fillmore EP. Thoracic wall and breast. In: Standring S, Tubbs RS. *Gray's anatomy: the anatomical basis of clinical practice.* 43rd ed. Philadelphia, PA: Elsevier Limited; 2026. Chapter 56, 1067-1090.e16
9. Fokin AA, Robicsek F. Poland's syndrome revisited. *Ann Thorac Surg.* 2002 Dec;74(6):2218-25.
10. Fonkalsrud EW, Anselmo DM. Less extensive techniques for repair of pectus carinatum: the undertreated chest deformity. *J Am Coll Surg.* 2004 Jun;198(6):898-905.
11. Fonkalsrud EW, Mendoza J. Open repair of pectus excavatum and carinatum deformities with minimal cartilage resection. *Am J Surg.* 2006 Jun;191(6):779-84.
12. Fonkalsrud EW, Salman T, Guo W, Gregg JP. Repair of pectus deformities with sternal support. *J Thorac Cardiovasc Surg.* 1994;107:37-42.
13. Gottlieb LJ, Reid RR, Slidell MB. Pediatric chest and trunk defects In: Rodriguez ED, Losee JE, Neligan PC, editors. *Plastic Surgery: Volume 3: Craniofacial, Head and Neck Surgery and Pediatric Plastic Surgery.* 4th ed. Elsevier; 2018. 888-911.e4.
14. Graham JM, Sanchez-Lara PA. Pectus excavatum and pectus carinatum. In: Graham JM, Sanchez-Lara PA. *Smith's recognizable patterns of human deformation.* 5th edition. Philadelphia, PA: Elsevier; 2025. Chapter 19, 145-153.

15. Graves CE, Hirose S, Raff GW, Iqbal CW, Imamura-Ching J, Christensen D, Fechter R, Kwiat D, Harrison MR. Magnetic Mini-Mover Procedure for pectus excavatum IV: FDA sponsored multicenter trial. *J Pediatr Surg*. 2017 Jun;52(6):913-919.
16. Guntheroth WG; Spiers PS. Cardiac function before and after surgery for pectus excavatum. *Am J Cardiol*. 2007 Jun;99(12):1762-4.
17. Haller JA Jr., Loughlin GM. Cardiorespiratory function is significantly improved following corrective surgery for severe pectus excavatum: proposed treatment guidelines. *J Cardiovasc Surg (Torino)*. 2000 Feb;41(1):125-30.
18. Harrison MR, Estefan-Ventura D, Fechter R, Moran AM Jr, Christensen D. Magnetic Mini-Mover Procedure for pectus excavatum: I. Development, design, and simulations for feasibility and safety. *J Pediatr Surg*. 2007 Jan;42(1):81-5; discussion 85-6.
19. Harrison MR, Gonzales KD, Bratton BJ, Christensen D, Curran PF, Fechter R, Hirose S. Magnetic mini-mover procedure for pectus excavatum III: safety and efficacy in a Food and Drug Administration-sponsored clinical trial. *J Pediatr Surg*. 2012 Jan;47(1):154-9.
20. Jaroszewski DE, Fonkalsrud EW. Repair of pectus chest deformities in 320 adult patients: 21 year experience. *Ann Thorac Surg*. 2007 Aug;84(2):429-33.
21. Johnson JN, Hartman TK, Pianosi PT, Driscoll DJ. Cardiorespiratory function after operation for pectus excavatum. *J Pediatr*. 2008 Sep;153(3):359-64.
22. Kelly RE Jr, Mellins RB, Shamberger RC, Mitchell KK, Lawson ML, Oldham KT, Azizkhan RG, Hebra AV, Nuss D, Goretsky MJ, Sharp RJ, Holcomb GW 3rd, Shim WK, Megison SM, Moss RL, Fecteau AH, Colombani PM, Cooper D, Bagley T, Quinn A, Moskowitz AB, Paulson JF. Multicenter study of pectus excavatum, final report: complications, static/exercise pulmonary function, and anatomic outcomes. *J Am Coll Surg*. 2013 Dec;217(6):1080-9.
23. Khan MI, Sun SX, Thompson Mancino A, Rosenbloom AL, Klimberg VS, Bland KI, Kass RB, Aydi ZB. Breast Physiology: normal and abnormal development and function. In: Klimberg VS, Gradishar WJ, Bland KI, Korourian S, White J, Copeland EM. *Bland and copeland's the breast*. 6th ed. Philadelphia, PA: Elsevier Inc. 2024. 5, 78-96.
24. Kubiak R, Habelt S, Hammer J, Häcker FM, Mayr J, Bielek J. Pulmonary function following completion of Minimally Invasive Repair for Pectus Excavatum (MIRPE). *Eur J Pediatr Surg*. 2007 Aug;17(4):255-60.
25. Lawson ML, Mellins RB, Tabangin M, Kelly Jr RE, Croitoru DP, Goretsky MJ, Nuss D. Impact of pectus excavatum on pulmonary function before and after repair with the Nuss procedure. *J Pediatr Surg*. 2005 Jan;40(1):174-80;discussion 180.
26. Lovegreen MT, Mogensen L. Congenital and acquired disorders. In: Murphy DP, Webster JB, Lovegreen W. *Atlas of orthoses and assistive devices*. 6th ed. Philadelphia, PA: Elsevier; 2026. 31, 326-333.e1.
27. Malek MH, Berger DE, Housh TJ, Marelich WD, Coburn JW, Beck TW. Cardiovascular function following surgical repair of pectus excavatum: a metaanalysis. *Chest*. 2006a Aug;130(2):506-16.

28. Malek MH, Berger DE, Marelich WD, Coburn JW, Beck TW, Housh TJ. Pulmonary function following surgical repair of pectus excavatum: a meta-analysis. *Eur J Cardiothorac Surg*. 2006b Oct;30(4):637-43.
29. Mao YZ, Tang S, Li S. Comparison of the Nuss versus Ravitch procedure for pectus excavatum repair: an updated meta-analysis. *J Pediatr Surg*. 2017 Oct;52(10):1545-1552.
30. Mourisse J, van der Heide S. Repair of Pectus Excavatum. In: Cohen E, editor. *Cohen's Comprehensive Thoracic Anesthesia*. Philadelphia, PA: Elsevier; 2022. 597-608.
31. Muthialu N, Kanani M, Smith OJ, Pearl R, Withey S. Chest wall reconstruction. In: Farhadieh RD, Bulstrode NW, Mehrara BJ, Cugno S. *Plastic surgery: principles and practice*. Philadelphia, PA: Elsevier Inc. 2022. 39, 599-609.
32. Pitt JB, Zeineddin S, Carter M, Perez A, Sands L, Kujawa S, Reynolds M, Abdullah F, Goldstein SD. Demographics of Anterior Chest Wall Deformity Patients: A Tertiary Children's Hospital Experience. *J Surg Res*. 2024 Jan;293:451-457. doi: 10.1016/j.jss.2023.09.040. Epub 2023 Oct 10. PMID: 37827024.
33. Poston PM, Patel SS, Rajput M, Rossi NO, Ghanamah MS, Davis JE, Turek JW. The correction index: setting the standard for recommending operative repair of pectus excavatum. *Ann Thorac Surg*. 2014 Apr;97(4):1176-9; discussion 1179-80.
34. Rodríguez-Granillo GA, Martínez-Ferro M, Capuñay C, Bellia-Munzón G, Deviggiano A, Raggio I, Fernandez-Rostello E, Hurtado Hoyo E, Bastarrika G, Carrascosa P. Preoperative multimodality imaging of pectus excavatum: State of the art review and call for standardization. *Eur J Radiol*. 2019 Aug;117:140-148.
35. Rust G, Westney GE. Pulmonary Medicine. In: Rakel RE, Rakel DP, editors. *Textbook of Family Medicine*. 9th ed. Philadelphia, PA: Elsevier Saunders; 2016. 236-273.e5
36. Sonaglioni A, Bruno A, Polymeropoulos A, Nicolosi GL, Lombardo M, Muti P. Prevalence of Mitral Valve Prolapse Among Individuals with Pectus Excavatum: A Systematic Review and Meta-Analysis. *Diagnostics (Basel)*. 2024 Nov 7;14(22):2488.
37. Sonaglioni A, Nicolosi GL, Trevisan R, Lombardo M, Grasso E, Gensini GF, Ambrosio G. The influence of pectus excavatum on cardiac kinetics and function in otherwise healthy individuals: A systematic review. *Int J Cardiol*. 2023 Jun 15;381:135-144.
38. St. Peter SD, Juang D, Garey CL, Laituri CA, Ostlie DJ, Sharp RJ, Snyder CL. A novel measure for pectus excavatum: the correction index. *J Pediatr Surg*. 2011 Dec;46(12):2270-3.
39. Sujka JA, St Peter SD. Quantification of pectus excavatum: Anatomic indices. *Semin Pediatr Surg*. 2018 Jun;27(3):122-126.
40. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. 510(k) Premarket Notification Database. Product code: HRS. Accessed Feb 2, 2026. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>

41. Zens TJ, Casar Berazaluce AM, Jenkins TM, Hardie W, Alsaied T, Tretter JT, Moore R, Foster K, Fleck RJ, Hanke RE, Colvin BE, Garrison AP, Kraemer A, Crotty E, Taylor M, Garcia VF, Brown RL. The Severity of Pectus Excavatum Defect Is Associated With Impaired Cardiopulmonary Function. Ann Thorac Surg. 2022 Sep;114(3):1015-1021.

42. Zganjer M, Zganjer V. Surgical correction of the funnel chest deformity in children. Int Orthop. 2010 Dec

Revision Details

Type of Revision	Summary of Changes	Date
Annual Review	<ul style="list-style-type: none">No clinical policy statement changes.	4/15/2026
Annual Review	<ul style="list-style-type: none">No clinical policy statement changes.	4/15/2025
Annual Review	<ul style="list-style-type: none">No clinical policy statement changes.	4/15/2024

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2026 The Cigna Group.