



Medical Coverage Policy

Effective Date1/15/2026

Next Review Date1/15/2027

Coverage Policy Number..... 0536

Prosthetic Devices

Table of Contents

Overview 2
 Coverage Policy 2
 Coding Information 5
 General Background..... 8
 Health Equity Considerations..... 16
 Medicare Coverage Determinations 17
 Appendix 17
 References..... 19
 Revision Details..... 25

Related Coverage Resources

- [Breast Reconstruction Following Mastectomy or Lumpectomy](#)
- [Gender Dysphoria Treatment](#)
- [Intraocular Lens Implant](#)
- [Male Sexual Dysfunction Treatment: Non-pharmacologic](#)
- [Orthotic Devices and Shoes](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer’s particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer’s benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer’s benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see “Coding Information” below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted

for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses prosthetic devices—fabricated items designed as replacements for missing body parts.

For information regarding prosthetic devices not addressed in this policy, please reference the applicable Cigna Medical Coverage Policy in the Related Coverage Resources section above.

Coverage Policy

Coverage for prosthetic devices varies across plans. Please refer to the customer's benefit plan document to determine benefit availability and the terms and conditions of coverage.

Microprocessor-controlled/computer-controlled/myoelectric devices are considered a type of power enhancement/controlled device.

GENERAL CRITERIA FOR A PROSTHETIC DEVICE

Functional Levels

Medical necessity for a lower limb prosthetic appliance is based on an individual's functional ability when using the prosthetic device. Functional ability is based on the following classification levels:

- **Level 0:** Does not have the ability or potential to ambulate or transfer safely with or without assistance and prosthesis does not enhance his/her quality of life or mobility.
- **Level 1:** Has the ability or potential to use prosthesis for transfers or ambulating on level surfaces at fixed cadence; typical of the limited and unlimited household ambulator.
- **Level 2:** Has the ability or potential for ambulating with the ability to traverse environmental barriers such as curbs, stairs or uneven surfaces; typical of the limited community ambulator.
- **Level 3:** Has the ability or potential for ambulating with variable cadence; typical of the community ambulator who has the ability to traverse most environmental barriers and may have vocational, therapeutic, or exercise activity that demands prosthetic utilization beyond simple locomotion.
- **Level 4:** Has the ability or potential for prosthetic ambulating that exceeds basic ambulating skills, exhibiting high impact, stress, or energy levels; typical of the prosthetic demands of the child, active adult, or athlete.

The following prosthetic devices are considered medically necessary when used to replace a missing or nonfunctional body part and when applicable medical necessity criteria listed below is met (Please note: prior authorization requirements may apply):

- External facial (e.g., nose, ear, midfacial, orbital, upper facial, hemifacial)
- Eye prosthesis (e.g., internal ocular, scleral shell)
- Lower extremity (e.g., foot, ankle, above/below knee)
- Upper extremity (e.g., finger, hand, wrist, above/below elbow, shoulder)
- Terminal devices, such as hands or hooks

Accessories to a prosthetic device are considered medically necessary when the accessory is required for the effective use of the prosthesis.

Not Medically Necessary

The following prosthetic devices are each considered not medically necessary:

- a lower limb prosthetic device for functional level 0
- additions/components that are not required for the effective use of the device
- prosthetic devices or additions/components not required for participation in normal activities of daily living, including those that are chiefly for convenience, for participation in recreational activities, or that otherwise exceed the medical needs of the individual (e.g., back-up/duplicate prosthetic devices, waterproof leg prosthesis [e.g., The Fin, used for swimming])

IRIS PROSTHESIS

An iris prosthesis (CPT® code 66683; HCPCS code C1839) is considered experimental, investigational or unproven for any indication, including but not limited to the treatment of full or partial aniridia.

EXTERNAL FACIAL PROSTHESIS

An external facial prosthesis is considered medically necessary when the prosthesis is prescribed to compensate for the loss or absence of facial tissue as a result of disease, injury, surgery, or congenital defect.

A duplicate external facial prosthesis is considered a convenience item and is considered not medically necessary.

UPPER LIMB: MYOELECTRIC PROSTHETIC DEVICE

If a benefit is available for an upper limb myoelectric device, the following medical necessity criteria apply.

An upper limb myoelectric prosthetic device is considered medically necessary for an individual with an amputation or congenital absence of a portion of an arm (e.g., hand, forearm, elbow) when ALL of the following criteria are met:

- The individual has sufficient cognitive ability to successfully utilize a myoelectric prosthetic device.
- The remaining musculature of the arm(s) contains the minimum microvolt threshold to allow operation of a myoelectric prosthetic device.

- A standard body-powered prosthetic device cannot be used or is insufficient to meet the functional needs of the individual in performing activities of daily living

An intent decoding module or pattern recognition add-on module for an upper limb myoelectric prosthetic device (e.g., Coapt Complete Control Gen2; Ottobock Myo Plus) (HCPCS code L6700) is considered experimental, investigational or unproven.

LOWER LIMB: MECHANICAL (NON-POWERED, NON-MICROPROCESSOR)

A single axis, fluid swing and stance phase control lower limb addition (HCPCS code L5828) is considered medically necessary when the individual is functional level 3 or greater.

LOWER LIMB: MICROPROCESSOR-CONTROLLED PROSTHETIC DEVICE

If a benefit is available for a microprocessor-controlled/computer-controlled lower limb prosthetic, the following medical necessity criteria apply.

Any of the following microprocessor-controlled prosthetics, including additions/components that are required for the effective use of the device (and consistent with the user's functional level), are considered medically necessary when the individual is functional level 3 or greater:

- a microprocessor-controlled ankle-foot prosthetic (HCPCS code L5973) for a transtibial amputee (i.e., below-the-knee)
- a microprocessor-controlled knee prosthetic (HCPCS code L5856) for a knee disarticulation amputee or a transfemoral amputee (i.e., above-the-knee)
- a combination microprocessor-controlled prosthetic/system (e.g., Linx), when a microprocessor-controlled prosthetic knee alone is inadequate to meet the functional needs of the individual (e.g., continued knee/foot instability due to environmental/anatomical barriers)

A microprocessor-controlled prosthetic is considered not medically necessary for any other indication.

LOWER LIMB: POWERED MICROPROCESSOR-CONTROLLED PROSTHETIC DEVICE

If a benefit is available for a powered or power-enhanced lower limb prosthetic, the following medical necessity criteria apply.

An endoskeletal knee-shin system (addition to a lower limb device) with powered and programmable flexion/extension assist control, including any type of motor(s) (HCPCS code L5859) (e.g., Össur Power Knee™) is considered medically necessary when ALL of the following criteria have been met:

- The individual has a swing and stance phase-type microprocessor controlled (electronic) knee (HCPCS code L5856).
- The individual is functional level 3 (K3) only*.

- The individual has a documented comorbidity of the spine and/or sound limb affecting hip extension and/or quadriceps function that impairs K3 level function with the use of a microprocessor-controlled knee alone.

***Note:** Coverage of this device is limited to individuals who are Functional Level 3; the device is not intended for high impact activity, sports, excessive loading, or heavy duty use.

The following powered prosthetic devices are each considered not medically necessary:

- a microprocessor-controlled ankle foot prosthetic with power assist (e.g., Ottobock Empower [HCPCS codes L5973, L5969])
- a powered lower limb prosthetic for any other indication

LOWER LIMB: VACUUM SUSPENSION SYSTEM

A vacuum suspension system (e.g., vacuum-assisted socket system [VASS]) (HCPCS code L5781) is considered medically necessary to control residual limb volume when there is contraindication to or failure of other socket-suspension systems (e.g., mechanical, passive suction) to adequately secure the limb to the prosthesis.

Coding Information

Notes:

1. This list of codes may not be all-inclusive since the American Medical Association (AMA) and Centers for Medicare and Medicaid Services (CMS) code updates may occur more frequently than policy updates.
2. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

IRIS PROSTHESIS

Considered Experimental/Investigational/Unproven:

CPT®* Codes	Description
66683	Implantation of iris prosthesis, including suture fixation and repair or removal of iris, when performed

HCPCS Codes	Description
C1839	Iris prosthesis

EXTERNAL FACIAL PROSTHESIS

Considered Medically Necessary when criteria in the applicable policy statements listed above are met and only when coverage is available under the plan for the specific device/component/item:

CPT®* Codes	Description
21088	Impression and custom preparation; facial prosthesis

UPPER LIMB: MYOELECTRIC PROSTHETIC DEVICE

Considered Medically Necessary when criteria in the applicable policy statements listed above are met and only when coverage is available under the plan for the specific device/component/item:

HCPCS Codes	Description
L6880	Electric hand, switch, or myoelectric controlled, independently articulating digits, any grasp pattern or combination of grasp patterns, includes motor(s)
L6881	Automatic grasp feature, addition to upper limb electric prosthetic terminal device
L6882	Microprocessor control feature, addition to upper limb prosthetic terminal device
L6935	Below elbow, external power, self-suspended inner socket, removable forearm shell, Otto Bock or equal electrodes, cables, two batteries and one charger, myoelectronic control of terminal device
L6955	Above elbow, external power, molded inner socket, removable humeral shell, internal locking elbow, forearm, Otto Bock or equal electrodes, cables, two batteries and one charger, myoelectronic control of terminal device
L7007	Electric hand, switch or myoelectric controlled, adult
L7259	Electronic wrist rotator, any type

Considered Experimental/Investigational/Unproven:

HCPCS Codes	Description
L6700	Upper extremity addition, external powered feature, myoelectronic control module, additional EMG inputs, pattern-recognition decoding intent movement

LOWER LIMB: MECHANICAL (NON-POWERED, NON-MICROPROCESSOR)

Considered Medically Necessary when used to report a component or addition to a lower limb prosthetic device when criteria in the applicable policy statements listed above are met and when coverage is available under the plan for the specific device/component/item:

HCPCS Codes	Description
L5828 [†]	Addition, endoskeletal knee-shin system, single axis, fluid swing and stance phase control
L5999 ^{††}	Lower extremity prosthesis, not otherwise specified

[†]Note: Considered medically necessary for functional level 3 or above.

^{††}Note: Considered medically necessary when used to report a medically necessary component or addition to a lower limb prosthetic device in the absence of a more specific code.

LOWER LIMB: MICROPROCESSOR-CONTROLLED PROSTHETIC DEVICES

Considered Medically Necessary when criteria in the applicable policy statements listed above are met and when benefits are available under the plan for a microprocessor-controlled prosthetic:

HCPCS Codes	Description
L5827	Endoskeletal knee-shin system, single axis, electromechanical swing and stance phase control, with or without shock absorption and stance extension damping
L5856	Addition to lower extremity prosthesis, endoskeletal knee-shin system, microprocessor control feature, swing and stance phase, includes electronic sensor(s), any type
L5973	Endoskeletal ankle foot system, microprocessor controlled feature, dorsiflexion and/or plantar flexion control, includes power source

Additional Components/Features of Microprocessor-Controlled Prosthetic Devices

Considered Medically Necessary when criteria in the applicable policy statements listed above are met and when benefits are available under the plan for a microprocessor-controlled prosthetic:

HCPCS Codes	Description
L5828	Addition, endoskeletal knee-shin system, single axis, fluid swing and stance phase control
L5999 [†]	Lower extremity prosthesis, not otherwise specified

[†]Note: Covered when used to report a medically necessary component/feature or addition to a lower limb prosthetic microprocessor-controlled device in the absence of a specific code.

LOWER LIMB: POWERED MICROPROCESSOR-CONTROLLED PROSTHETIC DEVICES

Considered Medically Necessary and when benefits are available for a power-controlled or power- assisted lower limb knee device (e.g., Ossur Power Knee):

HCPCS Codes	Description
L5859 [†]	Addition to lower extremity prosthesis, endoskeletal knee-shin system, powered and programmable flexion/extension assist control, includes any type motor(s)

[†]Note: L5859 requires K-3 functional level; the device is not intended for high impact activity, sports, excessive loading or heavy duty use.

Microprocessor-Controlled Ankle Foot Prosthetic with Power Assist (e.g., Ottobock Empower)

Considered Not Medically Necessary:

HCPCS Codes	Description
L5969	Addition, endoskeletal ankle-foot or ankle system, power assist, includes any type motor(s)
L5973	Endoskeletal ankle foot system, microprocessor controlled feature, dorsiflexion and/or plantar flexion control, includes power source

Additional Components/Features of Powered Prosthetic Devices, Including Power Assist Features

Considered Not Medically Necessary when reported in addition to a non-covered power-controlled (L5859, L5973) or power-assisted (L5969) prosthetic device:

HCPCS Codes	Description
L5828	Addition, endoskeletal knee-shin system, single axis, fluid swing and stance phase control
L5856	Addition to lower extremity prosthesis, endoskeletal knee-shin system, microprocessor control feature, swing and stance phase, includes electronic sensor(s), any type
L5969	Addition, endoskeletal ankle-foot or ankle system, power assist, includes any type motor(s)

LOWER LIMB: VACUUM SUSPENSION SYSTEM

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

HCPCS Codes	Description
L5781	Addition to lower limb prosthesis, vacuum pump, residual limb volume management and moisture evacuation system

***Current Procedural Terminology (CPT®) ©2025 American Medical Association: Chicago, IL.**

General Background

Prosthetic Devices

A prosthesis is an artificial device used to replace a missing body part and is intended to restore normal function. Prosthetic devices are secured or retained in place by harnesses or belts, by suction, or using anatomical structures; some devices such as facial prosthetics are held in place with the use of a skin adhesive. Additionally, devices may be held in place by implants, such as bone integrated titanium implants.

The following services and items are typically included in the allowance for a prosthetic device:

- the evaluation and fitting of the prosthesis
- the cost of base component parts and labor, as described in HCPCS base codes
- the repairs due to normal wear and tear during the 90-day period following the date of delivery

- adjustments of the prosthesis or the prosthetic component made when fitting the prosthesis or component and for 90 days from the date of delivery, when the adjustments are not necessitated by changes in the underlying tissue or the patient's functional ability

Iris Prostheses

An iris prosthesis is an implanted device proposed for the treatment of partial or complete aniridia. Aniridia is absence of the iris and may be associated with visual conditions such as glare, photophobia, glaucoma, corneal opacification, and/or cataract formation. The degree of vision loss varies. Treatment generally consists of contact lenses with iris prints and tinted eyeglasses. Various surgical techniques may also be used to repair iris defects, depending on the nature, extent and size of the defect (Ferro Desideri, et al., 2024).

The prosthetic iris device is made of foldable medical grade silicone which is then custom-sized and colored for each individual. The iris prosthetic is implanted surgically through a small incision, it is then unfolded, the edges are smoothed out and it is then held in place by anatomical structure of the eye or using sutures. It may be placed in the ciliary sulcus without sutures when there is a pre-existing intraocular lens, implanted into the capsular bag with a new intraocular lens, or can be sutured to the sclera, with or without an intraocular lens implant (IOL). The device purportedly reduces sensitivity to light while improving the appearance of the eye and visual acuity. Implant insertion can be done alone or in combination with cataract or lens fixation surgery.

U.S. Food and Drug Administration (FDA): The CustomFlex™ Artificial Iris (Clinical Research Consultants, Inc., [HumanOptics]) received premarket approval (PMA number P170039) by the FDA in May 2018 as an artificial iris intended for use in children and adults for the treatment of full or partial aniridia resulting from congenital aniridia, acquired defects, or other conditions associated with full or partial aniridia. The device is available with or without embedded fiber mesh for implantation, and may or may not be sutured. The FDA is requiring a post approval study to evaluate long term safety outcomes up to three years postoperatively for adults and five years for pediatric subjects.

Other implants have been investigated in the medical literature, however these devices have not been cleared or approved by the FDA (e.g., BrightOcular implants, used for cosmetic purposes). Some of the cosmetic devices have been associated with a high incidence of serious complications such as corneal decompensation, glaucoma, native iris trauma, intraocular inflammation, and cataract development, which may result in permanent structural damage or visual impairment (Ghaffari, 2021).

Literature Review: While there is a growing body of evidence in the peer-reviewed scientific literature evaluating use of the artificial iris, sample populations are small, lack a comparator group, study populations are heterogeneous, and surgical techniques vary, precluding generalization of overall safety and efficacy (Mayer, et al., 2021; Bonnet and Miller, 2020; Figueiredo and Snyder, 2020; Mayer, et al., 2019; Yoeruek and Bartz-Schmidt, 2019; Mayer, et al., 2018; Mostafa, et al., 2018; Mayer, et al., 2016; Rickmann, et al., 2016; Spitzer, et al., 2016). Several authors have reported high complications rates, both intra and post-operatively. As a result, strong evidence-based conclusions regarding safety and efficacy cannot be made. Additional clinical studies with longer follow-ups are needed to evaluate use of the device and impact on health outcomes.

Ayers et al. (2022) reported the results of a prospective, nonrandomized trial evaluating safety and efficacy of the CustomFlex Artificial Iris for treatment of partial or complete, congenital or acquired iris defects of various causes. Inclusion criteria were 22 years of age or greater, congenital or acquired iris defect and photophobia, glare sensitivity, or both, and pseudophakia,

phakia, or cataract in the study eye. The initial cohort involved 180 subjects, afterwards eligible adults were enrolled in a continued access cohort until the device received premarket approval from the FDA. Following at least four weeks post initial eye implantation fellow eye implantation was performed in 28 subjects. A compassionate use cohort (n=89) was also followed as part of the study protocol for individuals who did not meet one or more of the inclusion criteria. The authors reported subjects were reexamined one day following surgery and one week, one, three, six and 12 months after surgery. Three different techniques were used: (1) passive fixation within the capsular bag, (2) passive fixation within the ciliary sulcus, and (3) active suture fixation to residual iris tissue, the sclera, or an intraocular lens (IOL) that, in turn, was sutured to the sclera. Primary efficacy outcomes included a decrease in the severity of patient-reported photosensitivity (i.e., daytime and nighttime light sensitivity and daytime and nighttime glare), improvement in health-related quality of life, and improvement in postoperative cosmesis. Primary safety outcomes included cumulative IOL-related adverse events, cumulative surgery-related adverse events, and device-related adverse events. Secondary safety outcomes were tabulated and reported at the various study intervals and included changes in vision (CDVA, uncorrected distance visual acuity [UDVA], and manifest refraction), intraocular pressure (IOP), endothelial cell density (ECD), and slit-lamp observations. Endothelial cell density was measured at the screening visit and at six and 12 months after surgery if no corneal scarring, edema, or other pathologic features precluding measurement were present and was recorded as the average of three measurements obtained by noncontact specular or confocal microscopy. Results demonstrated a 59.7% reduction in marked to severe daytime light sensitivity ($p < 0.0001$), a 41.5% reduction in marked to severe nighttime light sensitivity ($p < 0.0001$), a 53.1% reduction in marked to severe daytime glare ($p < 0.0001$), and a 48.5% reduction in severe nighttime glare ($p < 0.0001$). A 15.4 point total score improvement was demonstrated in vision-related quality of life as measured by the 25-item National Eye Institute Visual Function Questionnaire (NEI VFQ-25) ($p < 0.0001$), and 93.8% of participants rated an improvement in cosmesis on the Global Aesthetic Improvement Scale at the 12-month postoperative examination. There was no loss of CDVA of $>$ two lines related to the device. Median ECD loss was 5.3% at six months after surgery and 7.2% at 12 months after surgery. The authors concluded that the artificial iris surpassed all key safety end points and met all key efficacy end points. The study was limited by the short length of follow-up.

Professional Societies/Organizations: The National Institute for Health and Care Excellence (NICE) (United Kingdom) published interventional procedures guidance for artificial iris insertion as treatment for acquired aniridia (NICE, 2020a). NICE reviewed evidence consisting of one non-randomized comparative trial, seven case series, and one case report. The primary efficacy outcomes included reduction in symptoms of glare, improvement in visual acuity, quality of life and other patient-reported outcomes. Key safety outcomes included need for explantation, infection, worsening visual acuity, glaucoma, and implant displacement. Within this document NICE concluded the "evidence on the safety and efficacy of artificial iris implant insertion for acquired aniridia is limited in quantity and quality. Therefore, this procedure should only be used with special arrangements for clinical governance, consent, and audit or research." Similarly, for congenital aniridia NICE concluded "evidence on the safety and efficacy of artificial iris implant insertion for congenital aniridia is inadequate in quantity and quality. Therefore, this procedure should only be used in the context of research" (NICE, 2020b).

External Facial Prostheses

External facial prostheses are used to replace lost or absent facial tissue that is the result of disease, injury, surgery, or a congenital defect, or they may be considered an alternative to reconstructive surgery. An external device is usually made from silicone materials and requires frequent removal and cleaning while a surgically implanted prosthetic device is typically removed and cleaned less often. The function of the external prosthesis is to protect exposed tissues, cover exposed cavities, and restore physical appearance.

Common types of external facial prostheses include the following:

- Auricular (ear): Restores all or part of the ear, function includes directing sound into the auditory canal; supporting eyeglasses and acting as a hearing aide, if required.
- Nasal (nose): Restores all or part of the nose and may include the nasal septum; functions to direct airflow to the nasopharynx and may also provide support for eyeglasses.
- Midfacial (nose and adjacent tissues): Restores part or all of the nose and significant adjacent facial tissue/structures, does not include the orbit or any intraoral maxillary prosthesis; adjacent facial tissue/structures include one or more of the following: soft tissue of the cheek, upper lip, or forehead.
- Orbital (orbit/eyelids): Restores the eyelids and the hard and soft tissue of the orbit, may include the eyebrow; functions to house the artificial eye, does not include the ocular prosthesis.
- Upper facial (orbit and adjacent tissues): Restores the orbit, plus significant adjacent facial tissue/structures, does not include the nose, any intraoral maxillary prosthesis or ocular prosthesis; adjacent facial tissue/structures include soft tissue of the cheek(s) or forehead.
- Hemifacial (nose, orbit and adjacent tissues): Restores part or all of the nose, the orbit, and significant adjacent facial tissue/structures, does not include any intraoral maxillary prosthesis or ocular prosthesis.
- Partial facial prosthesis: Restores a portion of the face, does not specifically involve the nose, orbit or ear.
- Nasal septal prosthesis: Prosthesis that occludes a hole in the nasal septum, does not include superficial nasal tissue.

Prosthetic devices may be secured or retained in place by anatomical structures; however, in most cases the device is held in place with the use of a skin adhesive. Additionally, some devices may be held in place by implants, such as bone integrated titanium implants. The method chosen to secure the device and the type of device are usually dependent upon factors such as the degree of deformity, the person's ability to handle maintenance routines, the individual's occupation and lifestyle, and the availability of assistance when needed.

Upper Limb Myoelectric Prosthetic Devices

The conventional prosthetic appliance for replacement of an upper extremity, either below or above the elbow, is a body-powered prosthesis with a terminal hand or hook device. A myoelectric device functions by means of electrical impulses and operates on rechargeable batteries requiring external cables or harnesses. It is a prosthetic device used as an alternative to a passive or conventional body-powered device which enables an amputee to adjust the force of his/her grip and an ability to both open and close the hand voluntarily. Myoelectric devices may be recommended for amputees who are unable to use body-powered devices or who require improved grip function/motion for performance of daily activities. Adults or children with above- or below-the-elbow amputations may use the device effectively, although as a child grows the prosthesis may require multiple socket replacements for proper fit and function.

A hybrid prosthesis is a device that uses a combination of myoelectric and body-powered technology to enhance the amputee's overall functionality, depending on the level and location of amputation. A hybrid device is indicated for high level amputations, (i.e., at or above the elbow) and consists of a body-powered device to control shoulder and elbow movement and a myoelectric device to control hand and wrist motion, allowing control of two joints at one time.

The integration of intent decoding modules with prostheses is an evolving area of interest. Intent decoding modules (IDMs) are advanced components added to myoelectric upper limb prosthetic devices, which use machine learning (pattern recognition) algorithms to translate myoelectric/electromyographic (EMG) signals from residual muscles into corresponding movement

commands to the prosthetic device. An accompanying app may also be used for configuration and feedback purposes. The purported benefits of machine learning pattern recognition control over conventional (direct) prosthesis control include more natural, fluid movements for improved function and precision. Examples of such devices include Coapt’s Complete Control Gen2 add-on controller kit, and Ottobock’s Myo Plus system.

U.S. Food and Drug Administration (FDA): The FDA generally considers upper limb myoelectric prostheses to be Class I devices, and exempt from premarket notification requirements. Prosthetic intent decoding modules (IDMs) may be cleared via the 510(k) premarket notification process, as Class II devices. The given indication for IDMs is for use exclusively with external prosthetic fittings of the upper limbs.

Device or Product	Identifier	Manufacturer
Complete Control Gen2	K191083	Coapt
Myo Plus	K191179	Otto Bock

*FDA product codes: GXY, IQZ

Note: Coverage decisions are not based solely on FDA approval. Device or product names are provided for example purposes only. Their inclusion does not indicate endorsement or preference for any specific brand or model. This list is not intended to reflect all available products or technologies.

Literature Review—Upper Limb Myoelectric Prostheses: The overall body of evidence in the peer-reviewed literature supports the safety and clinical benefits of upper limb myoelectric prostheses. Studies have demonstrated improvements in function and range of motion among individuals using these devices. Patient selection criteria are not uniformly defined, and direct comparisons to body-powered or passive devices remain limited. However, despite these limitations, the published literature supports the clinical value of myoelectric prostheses in enhancing upper limb function (Bouwsema, et al., 2010).

Literature Review—Intent Decoding/Pattern Recognition Modules: Evidence in the scientific peer-reviewed literature is insufficient to establish the efficacy and overall clinical utility of intent decoding or pattern recognition modules for upper limb prosthetic devices. Evidence is primarily in the form of case studies, case series, and randomized trials with small sample sizes, heterogenous patient populations, and mixed outcomes (Simon, et al., 2023b; Simon, et al., 2019; Woodward and Hargrove, 2019; Hargrove, et al., 2018; Hargrove, et al., 2017; Kuiken, et al., 2016; Kuiken, et al., 2009).

Lower Limb Prostheses

Prior to being fitted with a lower limb prosthetic device, the individual must demonstrate specific functional levels. A functional level is defined as a measurement of the capacity and potential of the individual to accomplish his/her expected post-rehabilitation daily function.

The Medicare Functional Classification Level (or “K level”) rating system consists of five classification levels and is used to gauge an individual's rehabilitation potential for using a lower limb prosthesis. The characteristics of individuals classified as K0 through K4 include the following (Centers for Medicare & Medicaid Services [CMS], 2017):

- Level 0: Does not have the ability or potential to ambulate or transfer safely with or without assistance and a prosthesis does not enhance their quality of life or mobility.
- Level 1: Has the ability or potential to use a prosthesis for transfers or ambulation on level surfaces at fixed cadence, typical of the limited and unlimited household ambulator.

- Level 2: Has the ability or potential for ambulation with the ability to transverse low level environmental barriers such as curbs, stairs or uneven surfaces. This level is typical of the limited community ambulator.
- Level 3: Has the ability or potential for ambulation with variable cadence, typical of the community ambulator who has the ability to transverse most environmental barriers and may have vocational, therapeutic, or exercise activity that demands prosthetic utilization beyond simple locomotion.
- Level 4: Has the ability or potential for prosthetic ambulation that exceeds the basic ambulation skills, exhibiting high impact, stress or energy levels typical of the prosthetic demands of the child, active adult, or athlete.

Lower limb prosthetic devices may be preparatory or permanent. A preparatory device is a prosthesis made soon after an amputation (approximately four weeks) as a temporary method of retraining a person to walk and balance while shrinking the residual limb. A permanent prosthesis is recommended when an individual has used a prosthetic device full time for a period of six months and when the limb volume has stabilized to a point where the socket fit remains relatively consistent for two to three weeks.

Components and/or additions to a prosthesis may be supported; the determination of medical necessity is based on the person's functional ability and expected functional potential as defined by the prosthetist and the ordering physician. Appropriate documentation supporting medical necessity must accompany requests submitted for prosthetic components and/or additions. Customizing prosthetic devices with enhanced features is generally not supported if activities of daily living can be met with standard devices.

Accessories that are necessary for the effective use of the prosthetic device may also be considered medically necessary devices. Accessories that are not necessary for the effective use of the device are considered not medically necessary. While some prosthetic manufacturers offer devices with waterproof features, including devices that are submersible (e.g., the Fin [Eschen Prosthetics and Orthotics Labs] [used for swimming]; Genium X3 and X4 [Ottobock], [waterproof microprocessor-controlled knee prosthetic devices]), when used for recreational purposes these prosthetic accessories/devices are considered a convenience item and not medically necessary.

Lower Limb Microprocessor-Controlled Devices—Knee: Microprocessor-controlled knee prosthetics, or microprocessor knees (MPKs), are sensor-equipped devices. The sensor detects when the knee is in full extension and adjusts the swing phases automatically, allowing a more natural pattern of walking at variable speeds (passive powered device).

Multiple devices are available that use various degrees of computer technology to enhance the clinical function of the basic mechanical knee design; all microprocessor-controlled systems do not have identical features and functions. Some devices have swing phase only, stance phase only, or swing and stance phase. Some of the devices currently available include the Ottobock C-Leg[®], and the Orion3 SmartIP (Blatchford Limited). The Genium X3 and X4 microprocessor devices by Ottobock are waterproof and submersible. The Kenevo prosthetic knee (Ottobock) is a device that is recommended for users with low to moderate mobility (indoor ambulation, limited outdoor ambulation) and is purported to better support those who use a walker, cane, crutch or wheelchair device. According to the manufacturer this device is not indicated for walking speeds greater than 3 km/hour and has a supported feature for stand-to-sit and sit-to-stand, wheelchair mode, and for putting on the prosthesis while seated.

The purported advantages of a microprocessor-controlled above-the-knee (AKA) prosthesis include:

- reduced energy expenditure of the amputee

- improved ability to walk on uneven ground
- improved ability to climb and descend stairs
- increased walking distance

U.S. Food and Drug Administration (FDA): The FDA generally considers lower limb microprocessor-controlled prostheses to be Class II (special controls) devices, and thus these devices are exempt from premarket notification requirements.

Device or Product	Identifier	Manufacturer
C-LEG	K991590	Ottobock

*FDA product codes: ISW, ISY, KFX

Note: Coverage decisions are not based solely on FDA approval. Device or product names are provided for example purposes only. Their inclusion does not indicate endorsement or preference for any specific brand or model. This list is not intended to reflect all available products or technologies.

Literature Review: Evidence in the peer-reviewed literature demonstrates that, for select populations, microprocessor-controlled knee prostheses provide important clinical benefits compared to conventional devices. Evidence is primarily in the form of case series, randomized trials, and systematic reviews, with studies reporting reductions in energy consumption, improved physical function, and a more symmetrical gait pattern when compared to a conventional. Additional research indicates decreased fall risk, as well as reductions in hip movement and metabolic requirements, particularly at faster walking speeds. Evidence of benefit is strongest for individuals in good health without significant comorbidities, and current literature supports the effective use of these devices for select populations where enhanced mobility and safety are clinically appropriate (Herrin, et al., 2025; McGrath et al., 2022; Carse et al., 2021; Campbell et al., 2020; Kaufman et al., 2018; Lansade et al., 2018; Fuenzalida Squella, et al., 2017; Prinsen, et al, 2017; Aldridge Whitehead et al., 2014; Highsmith, et al., 2013; Wolf, et al., 2013; Bellmann, et al., 2010). The evidence for individuals with lower functional levels (i.e., less active or primarily indoor users; < functional level 3) remains insufficient.

Lower Limb Microprocessor-Controlled Devices—Ankle: To enhance the basic mechanical design and mimic the action of a biological ankle, researchers have applied microprocessor technology to prosthetic feet. Stair ambulation is limited in the transtibial amputee because of neutral and fixed ankle position. Newer prosthetic ankles which adjust for ankle angle during swing phase and identify sloping gradients and ascent or descent of stairs are under investigation. One such microprocessor-controlled ankle foot prosthesis is the Proprio Foot® (Ossur). The Proprio Foot is a quasi-passive ankle that can actively change the ankle angle in the unloaded swing phase as the result of microprocessor-control and sensor technology. The device is passive (without power) while in stance phase. According to the manufacturer, the proposed benefits of microprocessor-controlled ankle movements include the ability to identify slopes and stairs, when ascending or descending stairs the device automatically adapts ankle position to enable the next step; allows the user to place both feet behind their knees when rising from a chair; and automatically gives a toe-lift allowing sufficient ground clearance when walking. The device is designed to promote a more symmetrical and balanced gait and is intended for use by transtibial amputees engaging in low to moderate impact activities who are classified as level K3 (i.e., community ambulatory, with the ability or potential for ambulation with variable cadence). It is not suitable for sport and high impact activities.

Literature Review: Evidence in the published peer-reviewed scientific literature evaluating the use of microprocessor-controlled ankle foot devices consists mainly of small comparative trials, pilot studies and case series involving small sample populations (Colas-Ribas, et al., 2022; Ernst,

et al., 2022; Kim, et al., 2021; Struchkov and Buckley, 2016; Darter and Wilken, 2014; Delussu, et al., 2013). The evidence demonstrates some clinical advantages for use compared to conventional ankle foot prostheses for individuals who are functional level 3 or greater. These devices may improve slope and uneven terrain ambulation, allowing larger range of motion of the ankle when compared with other conventional devices.

Combination Microprocessor-Controlled Knee-Ankle/Foot Prostheses: Combination microprocessor prosthetics are available integrating both a microprocessor knee and the ankle/foot device (e.g., SYMBIONIC® LEG 3 [Ossur, Iceland] [no longer commercially available]; Linx Limb System [Endolite/Blatchford]). One device, the SYMBIONIC® LEG 3 is a prosthetic that combines a microprocessor knee with a powered microprocessor ankle with proactive ankle flexion. The device purportedly has a more powerful knee actuator and new kinematic sensors for improved stability, increased support with stance flexion, and more rapid, and consistent swing extension. For a transfemoral amputee, combining both types of prosthetic devices theoretically enables a more natural and symmetrical gait when ambulating, decreasing energy expenditure, and offering increased stability. The device is intended for use by individuals who are Functional Level 3 or 4. The Linx prosthetic system is intended for individuals who are Functional Level 3 or greater; according to the manufacturer this system is an integrated prosthetic utilizing a microprocessor-controlled system in addition to sensors and actuators which simultaneously controls the knee and foot.

Lower Limb Powered Microprocessor-Controlled Prosthetic Devices

Powered Knee Prostheses: Powered prosthetic devices that use signals from muscle activity in the remaining limb to bend and straighten the device remain under investigation. These devices utilize sensors and electronics to process data and control movement and power of the knee. Examples of this type of device include the Power Knee™, manufactured by Ossur. According to the manufacturer, the Power Knee is described as a motorized device which contains a rechargeable battery pack. It is designed to replace muscle activity of the quadriceps muscle and uses artificial proprioception with sensors in order to anticipate and respond with the appropriate movement required for stepping (active powered device).

In comparison to a passive prosthetic knee, including a microprocessor device, the manufacturer suggests a power knee offers advantages such as powered extension with standing, controlled resistance with descending, and active flexion and extension during walking. The device controls the transition from a bent knee to an extended knee, at heel strike supports the individual's full body weight, and can help lift above-knee amputees out of a chair to a standing position. It is suggested the device helps to maintain walking speeds, assists with upward motion (required for stairs and inclines), and learns and responds to gait patterns. With the initial use of the device a practitioner must program and align the knee. Once programming and alignment are complete, the user needs only to press the power button to use the device. The device is compatible with a variety of dynamic flex-foot feet, must be re-charged daily and is not intended for high impact activity, sports, excessive loading or heavy duty use.

Powered Ankle-Foot Prostheses: Similar to the powered knee device, powered foot-ankle prosthetic devices are currently being developed. Two such devices are the BiOM® Ankle (BionX Medical Technologies, [previously iWalk, Inc., Bedford, MA) and the Empower prosthetic foot (Ottobock). The BiOM device (previously referred to as Powerfoot One) uses a combination of processors, sensors, motors, and springs that allow the user a powered push-off with taking steps. Theoretically the device replaces the action of the foot, Achilles tendon and calf muscle to result in a near normalized gait for amputees and is intended for amputees that are functional level 3 or 4.

Literature Review—Powered Ankle-Foot Prostheses: There is limited evidence in the published peer-reviewed literature to demonstrate the safety, efficacy, and overall clinical utility of

powered ankle-foot prostheses compared to conventional devices. Most available research focuses on device design and biomechanical performance, with few comparative clinical trials. Although some studies report outcomes such as kinematic improvements, reduced energy expenditure, and enhanced biomechanical function with powered ankle-foot prostheses, these studies typically involve small sample sizes, short-term follow-ups, and lack control groups (Rabago, et al., 2016; Russel Esposito, et al., 2016; Ingraham, et al., 2016; Simon, et al., 2016; Gates, et al., 2013; Grabowski and D'Andrea, 2013; Takahashi and Stanhope, 2013; Aldridge, et al., 2012; Herr and Grabowski, 2012).

Lower Limb Vacuum Suspension Systems: Suspension systems for lower limb prostheses keep the prosthesis in place, ensuring a good fit between the socket and residual limb. The intended function is to provide a connection that reduces rotational and shearing forces which can result in skin breakdown as well provide for balance and steady gait. Various types of suspension systems are available and include those that are primarily mechanical or suction-type systems. Mechanical systems involve the use of belts, straps, or sleeves, for example, to attach the device to the residual limb. Suction-type systems function by way of a negative pressure created between the socket and insert/liner. These devices can be passive (air escapes while donning via a one-way valve) or active (suction pump evacuates the air). Passive systems involve the use of a soft liner, a one-way valve and a donning sleeve. A liner is placed over the limb, the limb is placed in the socket and the force of one's body weight upon standing expels excess air through the valve creating a seal. With active suction devices the sleeve creates a seal around the edge of the socket and a pump and exhaust remove the excess air between the socket and the liner to ensure a secure fit.

Various vacuum suction-type devices (mechanical or electrical) are available and include the Vacuum-Assisted Socket System (VASS) (Ottobock Harmony Vacuum-Assisted Socket System), and the LimbLogic prosthetic vacuum suspension system (WillowWood). Each device is a vacuum suction-type suspension system that manufacturers claim helps control volume fluctuation in the residual limbs of lower-extremity amputees, reduces forces to the limbs, and improves both suspension and proprioception without restricting vascular flow. Although patient selection criteria have not been firmly established, the device has been proposed for individuals with non-healing skin ulcerations located on the stump and/or when other socket systems have failed to provide a secure fit.

Literature Review: There is evidence to support vacuum systems decrease limb volume fluctuations, improve socket fit, reduce inside movement for some individuals, decrease falls, and improve comfort and satisfaction (Youngblood, et al., 2020; Rosenblatt and Ehrhardt, 2017; Darter, et al., 2016; Gholizadeh, et al., 2016; Hoskins, et al., 2014; Samitier, et al., 2014; Traballese, et al., 2012; Klute, et al., 2011; Beil, et al., 2002). For individuals where other types of suspension systems have failed to provide a secure fit or are contraindicated, a vacuum suction-type suspension system may be considered an effective alternative. The choice of a suspension system is determined by factors such as activity level, residual stump shape, age, and health status.

Health Equity Considerations

Health equity is the highest level of health for all people; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which people are born, grow, live, work, and age.

Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include safe housing,

transportation, and neighborhoods; racism, discrimination and violence; education, job opportunities and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

Medicare Coverage Determinations

	Contractor	Determination Name/Number	Revision Effective Date
NCD	National	No Determination found	
LCD	CGS Administrators; Noridian Healthcare Solutions	Lower Limb Prostheses (L33787)	10/1/2025
LCD	CGS Administrators; Noridian Healthcare Solutions	Facial Prostheses (L33738)	1/1/2020
LCD	CGS Administrators; Noridian Healthcare Solutions	Eye Prostheses (L33737)	1/1/2020

Note: Please review the current Medicare Policy for the most up-to-date information. (NCD = National Coverage Determination; LCD = Local Coverage Determination)

Appendix

Lower Limb Prosthetic “Device to Coding” Crosswalk

Please note, this list is for informational purposes only; it DOES NOT imply coverage or non-coverage of a device, or guarantee claim reimbursement. Coding may vary according to manufacturer.

Device Name	Brief Description	Manufacturer	HCPCS Code(s)
Allux™ 2	Microprocessor-controlled knee	Nabtesco	L5856, L5848, L5845, L5615
Avior	Microprocessor-controlled knee (for activity/mobility levels K1, K2)	Blatchford	L5856, L5850, L5848, L5845, L5615, L5925
Bio Leg®	Microprocessor-controlled motor-powered knee	BionicM	L5859, L5856, L5827
C-Leg	Microprocessor-controlled knee	Ottobock	L5856, L5850, L5848, L5845, L5828, L5925
Elan ElanIC	Microprocessor-controlled ankle foot (ElanIC is waterproof)	Blatchford	L5973

Device Name	Brief Description	Manufacturer	HCPCS Code(s)
Empower	Microprocessor-controlled ankle foot (power)	Ottobock	L5969, L5973
Genium Genium X3 Genium X4	Microprocessor-controlled knee (X3 and X4 are waterproof)	Ottobock	L5999
Icon	Microprocessor-controlled knee	College Park Industries	L5856, L5850, L5848, L5845, L5828, L5925
INTUY® Knee	Microprocessor-controlled motor-powered knee	WillowWood	L5859, L5856, L5827
Kenevo	Microprocessor-controlled knee	Ottobock	L5856, L5850, L5848, L5845, L5828, L5925
Kinnex™ 2.0	Microprocessor-controlled ankle foot	Proteor	L5973
Kneuro	Microprocessor-controlled knee (for activity/mobility levels K2-K4)	BrainRobotics	L5856, L5850, L5848, L5845, L5828, L5925
Linx	Combination microprocessor-controlled knee and foot	Blatchford	L5856, L5850, L5848, L5845, L5828, L5925, L5973
Meridium	Microprocessor-controlled ankle foot	Ottobock	L5973
Navii®	Microprocessor-controlled knee (for activity/mobility levels K2-K4; waterproof)	Össur	L5856, L5850, L5848, L5845, L5828, L5925
Odyssey® iQ	Microprocessor-controlled ankle foot	College Park Industries	L5973
Orion3	Microprocessor-controlled knee	Blatchford	L5856, L5850, L5848, L5845, L5828, L5925
Power Knee™	Microprocessor-controlled motor-powered knee	Össur	L5859, L5856, L5827
Plié 3®	Microprocessor-controlled knee (submersible)	Proteor	L5856, L5850, L5848, L5845, L5828
Pro-Flex® Pivot	Mechanical ankle foot with vacuum suspension	Össur	L5781, L5999
Proprio Foot®	Microprocessor-controlled ankle foot (for activity/mobility levels K2, K3; waterproof)	Össur	L5973
Quattro®	Microprocessor-controlled knee	Proteor	L5856, L5828, L5848, L5845, L5850, L5925

Device Name	Brief Description	Manufacturer	HCPCS Code(s)
Rheo Knee® Rheo Knee® XC	Microprocessor-controlled knee (Rheo Knee XC supports early rehabilitation to full recovery)	Össur	L5856, L5850, L5848, L5845, L5828, L5925
SLK™ 3.0	Microprocessor-controlled knee	DAW Industries	L5856, L5848, L5845, L5613
SmartIP	Microprocessor-controlled knee, with weight activated stance control	Blatchford	L5830, L5857 (+L5845 for Stanceflex models only)
SYNSYS	Combination microprocessor full-leg system (for activity/mobility levels K2, low to moderate K3)	Proteor	L5856, L5850, L5848, L5845, L5828, L5925, L5973

References

1. Aldridge JM, Sturdy JT, Wilken JM. Stair ascent kinematics and kinetics with a powered lower leg system following transtibial amputation. *Gait Posture*. 2012 Jun;36(2):291-5.
2. Aldridge Whitehead JM, Wolf EJ, Scoville CR, Wilken JM. Does a Microprocessor-controlled Prosthetic Knee Affect Stair Ascent Strategies in Persons With Transfemoral Amputation? *Clin Orthop Relat Res*. 2014 Feb 11.
3. Alzeer AM, Bhaskar Raj N, Shahine EM, Nadiah WA. Impacts of Microprocessor-Controlled Versus Non-microprocessor-Controlled Prosthetic Knee Joints Among Transfemoral Amputees on Functional Outcomes: A Comparative Study. *Cureus*. 2022 Apr 21;14(4):e24331.
4. Ang M, Tan D. Anterior segment reconstruction with artificial iris and Descemet membrane endothelial keratoplasty: a staged surgical approach. *Br J Ophthalmol*. 2021 Feb 26:bjophthalmol-2020-317906.
5. Ayres BD, Fant BS, Landis ZC, Miller KM, Stulting RD, Cionni RJ, Fram NR, Hamilton S, Hardten DR, Koch DD, Masket S, Price FW Jr, Rosenthal KJ, Hamill MB, Snyder ME. Results of the United States Food and Drug Administration Clinical Trial of the CustomFlex Artificial Iris. *Ophthalmology*. 2022 Jun;129(6):614-625.
6. Barnett CT, Hughes LD, Sullivan AE, Strutzenberger G, Levick JL, Bisele M, De Asha AR. Exploring the interaction of knee and ankle component use on mobility test performance in people with unilateral transfemoral amputation. *Prosthet Orthot Int*. 2021 Dec 1;45(6):470-476.
7. Beil TL, Street GM, Covey SJ. Interface pressures during ambulation using suction and vacuum-assisted prosthetic sockets. *J Rehabil Res Dev*. 2002 Nov-Dec;39(6):693-700

8. Bellmann M, Schmalz T, Blumentritt S. Comparative biomechanical analysis of current microprocessor-controlled prosthetic knee joints. *Arch Phys Med Rehabil.* 2010 Apr;91(4):644-52.
9. Blatchford. Products. Accessed Dec 5, 2025. Available at URL address: <https://www.blatchfordmobility.com/en-us/products/>
10. Blatchford. Suggested L Codes. Accessed Dec 5, 2025. Available at URL address: <https://www.blatchfordmobility.com/media/3cagpui0/203307228-website-l-codes-iss15-aw-int.pdf>
11. Bonnet C, Miller KM. Safety and efficacy of custom foldable silicone artificial iris implantation: prospective compassionate-use case series. *J Cataract Refract Surg.* 2020 Jun;46(6):893-901.
12. Bouwsema H, van der Sluis CK, Bongers RM. Learning to control opening and closing a myoelectric hand. *Arch Phys Med Rehabil.* 2010 Sep;91(9):1442-6.
13. Campbell JH, Stevens PM, Wurdeman SR. OASIS 1: Retrospective analysis of four different microprocessor knee types. *J Rehabil Assist Technol Eng.* 2020 Nov 5;7:2055668320968476.
14. Carse B, Scott H, Brady L, Colvin J. Evaluation of gait outcomes for individuals with established unilateral transfemoral amputation following the provision of microprocessor controlled knees in the context of a clinical service. *Prosthet Orthot Int.* 2021 Jun 1;45(3):254-261.
15. Centers for Medicare and Medicaid Services (CMS). Health Technology Assessment. Lower Limb Prosthetic Workgroup Consensus Document. Sep 2017. Accessed Dec 5, 2025. Available at URL address: https://www.cms.gov/Medicare/Coverage/DeterminationProcess/downloads/LLP_Consensus_Document.pdf
16. Centers for Medicare and Medicaid Services (CMS). Medicare Coverage Database. Accessed Dec 3, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/search.aspx>
17. Coapt. Technology. Accessed Dec 5, 2025. Available at URL address: <https://coaptengineering.com/technology>
18. Colas-Ribas C, Martinet N, Audat G, Bruneau A, Paysant J, Abraham P. Effects of a microprocessor-controlled ankle-foot unit on energy expenditure, quality of life, and postural stability in persons with transtibial amputation: An unblinded, randomized, controlled, cross-over study. *Prosthet Orthot Int.* 2022 Dec 1;46(6):541-548. doi: 10.1097/PXR.000000000000187. PMID: 36515900.
19. Crawford AZ, Freundlich SEN, Lim J, McGhee CNJ. Endocapsular artificial iris implantation for iris defects: Reducing symptoms, restoring visual function and improving cosmesis. *Clin Exp Ophthalmol.* 2022 Jul;50(5):490-499.
20. Darter BJ, Sinitski K, Wilken JM. Axial bone-socket displacement for persons with a traumatic transtibial amputation: The effect of elevated vacuum suspension at progressive body-weight loads. *Prosthet Orthot Int.* 2015 Sep 30.

21. Darter BJ, Wilken JM. Energetic consequences of using a prosthesis with adaptive ankle motion during slope walking in persons with a transtibial amputation. *Prosthet Orthot Int*. 2013 Mar 22.
22. Davie-Smith F, Carse B. Comparison of patient-reported and functional outcomes following transition from mechanical to microprocessor knee in the low-activity user with a unilateral transfemoral amputation. *Prosthet Orthot Int*. 2021 Jun 1;45(3):198-204.
23. Delussu AS, Brunelli S, Paradisi F, Iosa M, Pellegrini R, Zenardi D, Traballese M. Assessment of the effects of carbon fiber and bionic foot during overground and treadmill walking in transtibial amputees. *Gait Posture*. 2013 Sep;38(4):876-82.
24. Ernst M, Altenburg B, Schmalz T, Kannenberg A, Bellmann M. Benefits of a microprocessor-controlled prosthetic foot for ascending and descending slopes. *J Neuroeng Rehabil*. 2022 Jan 28;19(1):9.
25. Ferro Desideri L, Arun K, Doherty G, Bernardi E, Anguita R. Iris Reconstruction: A Surgeon's Guide. *J Clin Med*. 2024 May 4;13(9):2706.
26. Figueiredo GB, Snyder ME. Long-term follow-up of a custom-made prosthetic iris device in patients with congenital aniridia. *J Cataract Refract Surg*. 2020 Jun;46(6):879-887.
27. Fuenzalida Squella SA, Kannenberg A, Brandão Benetti Â. Enhancement of a prosthetic knee with a microprocessor-controlled gait phase switch reduces falls and improves balance confidence and gait speed in community ambulators with unilateral transfemoral amputation. *Prosthet Orthot Int*. 2017 Jul 1:309364617716207.
28. Gates DH, Aldridge JM, Wilken JM. Kinematic comparison of walking on uneven ground using powered and unpowered prostheses. *Clin Biomech (Bristol, Avon)*. 2013 Apr;28(4):467-72.
29. Ghaffari R, Aldave AJ, Al-Hashimi S, Miller KM. Complications of Cosmetic Artificial Iris Implantation and Post Explantation Outcomes. *Am J Ophthalmol*. 2021 Jun;226:156-164.
30. Gholizadeh H, Lemaire ED, Eshraghi A. The evidence-base for elevated vacuum in lower limb prosthetics: Literature review and professional feedback. *Clin Biomech (Bristol, Avon)*. 2016 Jun 22;37:108-116.
31. Grabowski AM, D'Andrea S. Effects of powered ankle-foot prosthesis on kinetic loading of the unaffected leg during level-ground walking. *Neuroeng Rehabil*. 2013 Jun 7;10:49.
32. Hafner BJ, Halsne EG, Morgan SJ, Morgenroth DC, Humbert AT. Effects of prosthetic feet on metabolic energy expenditure in people with transtibial amputation: A systematic review and meta-analysis. *PM R*. 2021 Aug 14.
33. Hahn A, Bueschges S, Prager M, Kannenberg A. The effect of microprocessor controlled exo-prosthetic knees on limited community ambulators: systematic review and meta-analysis. *Disabil Rehabil*. 2021 Oct 25:1-19.
34. Hargrove L, Miller L, Turner K, Kuiken T. Control within a virtual environment is correlated to functional outcomes when using a physical prosthesis. *J Neuroeng Rehabil*. 2018 Sep 5;15(Suppl 1):60.

35. Hargrove LJ, Miller LA, Turner K, Kuiken TA. Myoelectric Pattern Recognition Outperforms Direct Control for Transhumeral Amputees with Targeted Muscle Reinnervation: A Randomized Clinical Trial. *Sci Rep*. 2017 Oct 23;7(1):13840.
36. Herr HM, Grabowski AM. Bionic ankle-foot prosthesis normalizes walking gait for persons with leg amputation. *Proc Biol Sci*. 2012 Feb 7;279(1728):457-64.
37. Herrin K, Kestur S, Zhou S, O'Sullivan G, Snow T, Childers WL, Young A. Toward personalizing prosthesis prescription: A take-home study of three microprocessor-controlled prosthetic knees: A randomized crossover study. *PM R*. 2025 Oct 24. doi: 10.1002/pmrj.70028. Epub ahead of print. PMID: 41133928.
38. Highsmith MJ, Kahle JT, Miro RM, Mengelkoch LJ. Ramp descent performance with the C-Leg and interrater reliability of the Hill Assessment Index. *Prosthet Orthot Int* 2013 Jan 17.
39. Hoskins RD, Sutton EE, Kinor D, Schaeffer JM, Fatone S. Using vacuum-assisted suspension to manage residual limb wounds in persons with transtibial amputation: a case series. *Prosthet Orthot Int*. 2014 Feb;38(1):68-74.
40. Ingraham KA, Fey NP, Simon AM, Hargrove LJ. Assessing the Relative Contributions of Active Ankle and Knee Assistance to the Walking Mechanics of Transfemoral Amputees Using a Powered Prosthesis. *PLoS One*. 2016 Jan 25;11(1):e0147661.
41. Jakob-Girbig J, Salewsky S, Meller D. Use of the Ophtec Artificial Iris Model C1 in Patients with Aniridia/Aphakia. *Klin Monbl Augenheilkd*. 2021 Jul;238(7):803-807.
42. Jayaraman C, Mummidisetty CK, Albert MV, Lipschutz R, Hoppe-Ludwig S, Mathur G, Jayaraman A. Using a microprocessor knee (C-Leg) with appropriate foot transitioned individuals with dysvascular transfemoral amputations to higher performance levels: a longitudinal randomized clinical trial. *J Neuroeng Rehabil*. 2021 May 25;18(1):88.
43. Kaufman KR, Bernhardt KA, Symms K. Functional assessment and satisfaction of transfemoral amputees with low mobility (FASTK2): A clinical trial of microprocessor-controlled vs. non-microprocessor-controlled knees. *Clin Biomech (Bristol)*. 2018 Oct;58:116-122.
44. Kestur S, Zhou S, O'Sullivan G, Young A, Herrin K. Comparing the lower limb joint biomechanics of the Power Knee, C-Leg and Rheo Knee during ramp and stair ambulation. *J Biomech*. 2024 Jun;171:112201.
45. Kim J, Wensman J, Colabianchi N, Gates DH. The influence of powered prostheses on user perspectives, metabolics, and activity: a randomized crossover trial. *J Neuroeng Rehabil*. 2021 Mar 16;18(1):49.
46. Klute GK, Berge JS, Biggs W, Pongnumkul S, Popovic Z, Curless B. Vacuum-assisted socket suspension compared with pin suspension for lower extremity amputees: effect on fit, activity, and limb volume. *Arch Phys Med Rehabil*. 2011 Oct;92(10):1570-5.
47. Kuiken TA, Li G, Lock BA, Lipschutz RD, Miller LA, Stubblefield KA, Englehart KB. Targeted muscle reinnervation for real-time myoelectric control of multifunction artificial arms. *JAMA*. 2009 Feb 11;301(6):619-28.

48. Kuiken TA, Miller LA, Turner K, Hargrove LJ. A Comparison of Pattern Recognition Control and Direct Control of a Multiple Degree-of-Freedom Transradial Prosthesis. *IEEE J Transl Eng Health Med.* 2016 Nov 22;4:2100508.
49. Lansade C, Vicaut E, Paysant J, Ménager D, Cristina MC, Braatz F, Domayer S, Pérennou D, Chiesa G. Mobility and satisfaction with a microprocessor-controlled knee in moderately active amputees: A multi-centric randomized crossover trial. *Ann Phys Rehabil Med.* 2018 Sep;61(5):278-285.
50. Mayer CS, Baur ID, Storr J, Khoramnia R. Bilateral Artificial Iris implantation in patients with bilateral iris defects. *Am J Ophthalmol Case Rep.* 2021 Apr 30;22:101108.
51. Mayer CS, Laubichler AE, Khoramnia R, et al. Challenges and complication management in novel artificial iris Implantation. *J Ophthalmol.* 2018 Sep 23; 2018:3262068.
52. Mayer CS, Laubichler AE, Masyk M, et al. Residual iris retraction syndrome after artificial iris implantation. *Am J Ophthalmol.* 2019 Mar;199:159-166.
53. Mayer CS, Reznicek L, Hoffmann AE. Pupillary reconstruction and outcome after artificial iris implantation. *Ophthalmology.* 2016 May; 123(5):1011-1018.
54. Mostafa YS, Osman AA, Hassanein DH, et al. Iris reconstruction using artificial iris prosthesis for management of aniridia. *Eur J Ophthalmol.* 2018 Jan;28(1):103-107.
55. Müller R, Tronicke L, Abel R, Lechler K. Prosthetic push-off power in trans-tibial amputee level ground walking: A systematic review. *PLoS One.* 2019 Nov 19;14(11):e0225032. doi: 10.1371/journal.pone.0225032. PMID: 31743353; PMCID: PMC6863538.
56. National Institute for Health and Care Excellence (NICE). Artificial iris insertion for acquired aniridia. IPG674. Jul 22, 2020a. Accessed Dec 3, 2025. Available at URL address: <https://www.nice.org.uk/guidance/ipg674>
57. National Institute for Health and Care Excellence (NICE). Artificial iris insertion for congenital aniridia. IPG675. Published Jul 22, 2020b. Accessed Dec 3, 2025. Available at URL address: <https://www.nice.org.uk/guidance/ipg675>
58. Nissen K, Skibsted SP, Bagger MM, Ejstrup R, Faber C, Riise P, Heegaard S, Kiilgaard JF. Outcome of combined cataract surgery and iris prosthesis implantation in iris melanoma patients previously treated with iridocyclectomy: A national retrospective non-comparative case series. *Acta Ophthalmol.* 2024 Aug;102(5):581-589.
59. Össur. Prosthetics. Accessed Dec 5, 2025. Available at URL address: <https://www.ossur.com/en-us/prosthetics>
60. Ottobock. Myo Plus TR. Accessed Dec 5, 2025. Available at URL address: <https://www.ottobock.com/en-us/product/13E520-61077>
61. Ottobock. Prosthetics. Accessed Dec 5, 2025. Available at URL address: <https://shop.ottobock.us/Prosthetics/c/1>

62. Prinsen EC, Nederhand MJ, Sveinsdóttir HS, et al. The influence of a user-adaptive prosthetic knee across varying walking speeds: A randomized cross-over trial. *Gait Posture*. 2017 Jan;51:254-260.
63. Proteor. PDAC Verified Products. Accessed Dec 5, 2025. Available at URL address: <https://us.proteor.com/pdac-verified-products-demo/>
64. Rabago CA, Aldridge Whitehead J, Wilken JM. Evaluation of a Powered Ankle-Foot Prosthesis during Slope Ascent Gait. *PLoS One*. 2016 Dec 15;11(12):e0166815.
65. Rickmann A, Szurman P, Januschowski K, et al. Long-term results after artificial iris implantation in patients with aniridia. *Graefes Arch Clin Exp Ophthalmol*. 2016 Jun; 254(7):1419-1424.
66. Romano D, Bremond-Gignac D, Barbany M, Rahman A, Muring L, Semeraro F, Cursiefen C, Lagali N, Romano V. Artificial iris implantation in congenital aniridia A systematic review. *Surv Ophthalmol*. 2022 Nov 12:S0039-6257(22)00152-7.
67. Rosenblatt NJ, Ehrhardt T. The effect of vacuum assisted socket suspension on prospective, community-based falls by users of lower limb prostheses. *Gait Posture*. 2017 Jun;55:100-104. doi: 10.1016/j.gaitpost.2017.03.038. Epub 2017 Apr 8. PMID: 28433866.
68. Russell Esposito E, Aldridge Whitehead JM, Wilken JM. Step-to-step transition work during level and inclined walking using passive and powered ankle-foot prostheses. *Prosthet Orthot Int*. 2016 Jun;40(3):311-9.
69. Samitier CB, Guirao L, Costea M, Camós JM, Pleguezuelos E. The benefits of using a vacuum-assisted socket system to improve balance and gait in elderly transtibial amputees. *Prosthet Orthot Int*. 2014 Sep 26.
70. Simon AM, Fey NP, Ingraham KA, Finucane SB, Halsne EG, Hargrove LJ. Improved Weight-Bearing Symmetry for Transfemoral Amputees During Standing Up and Sitting Down With a Powered Knee-Ankle Prosthesis. *Arch Phys Med Rehabil*. 2016 Jul;97(7):1100-6.
71. Simon AM, Turner KL, Miller LA, Dumanian GA, Potter BK, Beachler MD, Hargrove LJ, Kuiken TA. Myoelectric prosthesis hand grasp control following targeted muscle reinnervation in individuals with transradial amputation. *PLoS One*. 2023a Jan 26;18(1):e0280210.
72. Simon AM, Turner KL, Miller LA, Hargrove LJ, Kuiken TA. Pattern recognition and direct control home use of a multi-articulating hand prosthesis. *IEEE Int Conf Rehabil Robot*. 2019 Jun;2019:386-391.
73. Simon AM, Turner KL, Miller LA, Potter BK, Beachler MD, Dumanian GA, Hargrove LJ, Kuiken TA. User Performance With a Transradial Multi-Articulating Hand Prosthesis During Pattern Recognition and Direct Control Home Use. *IEEE Trans Neural Syst Rehabil Eng*. 2023b;31:271-281.
74. Struchkov V, Buckley JG. Biomechanics of ramp descent in unilateral trans-tibial amputees: Comparison of a microprocessor controlled foot with conventional ankle-foot mechanisms. *Clin Biomech (Bristol, Avon)*. 2016 Feb;32:164-70.

75. Takahashi KZ, Stanhope SJ. Mechanical energy profiles of the combined ankle-foot system in normal gait: insights for prosthetic designs. *Gait Posture*. 2013 Sep;38(4):818-23. doi: 10.1016/j.gaitpost.2013.04.002. Epub 2013 Apr 28. PMID: 23628408.
76. Traballesi M, Delussu AS, Fusco A, Iosa M, Averna T, Pellegrini R, Brunelli S. Residual limb wounds or ulcers heal in transtibial amputees using an active suction socket system. A randomized controlled study. *Eur J Phys Rehabil Med*. 2012 May 28.
77. U.S. Food and Drug Administration (FDA). 510(k) premarket notification database. Product code(s): GXY, IQZ, ISW, ISY, KFX. Page Last Updated: Dec 1, 2025. Accessed Dec 3, 2025. Available at URL address: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>
78. U.S. Food Drug Administration (FDA). Premarket approval (PMA) database. Product code(s): QBT. Page Last Updated: Dec 1, 2025. Accessed Dec 3, 2025. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm>
79. Weber EL, Stevens PM, England DL, Swilley VD, Wurdeman SR. Microprocessor feet improve prosthetic mobility and physical function relative to non-microprocessor feet. *J Rehabil Assist Technol Eng*. 2022 Jul 11;9:20556683221113320.
80. Wolf EJ, Everding VQ, Linberg AA, Czerniecki JM, Gambel CJ. Comparison of the Power Knee and C-Leg during step-up and sit-to-stand tasks. *Gait Posture*. 2013 Jul;38(3):397-402.
81. Woodward RB, Hargrove LJ. Adapting myoelectric control in real-time using a virtual environment. *J Neuroeng Rehabil*. 2019 Jan 16;16(1):11.
82. Yoeruek E, Bartz-Schmidt KU. A new knotless technique for combined transscleral fixation of artificial iris, posterior chamber intraocular lens, and penetrating keratoplasty. *Eye (Lond)*. 2019 Mar;33(3):358-362.
83. Youngblood RT, Brzostowski JT, Hafner BJ, Czerniecki JM, Allyn KJ, Foster RL, Sanders JE. Effectiveness of elevated vacuum and suction prosthetic suspension systems in managing daily residual limb fluid volume change in people with transtibial amputation. *Prosthet Orthot Int*. 2020 Jun;44(3):155-163.

Revision Details

Type of Revision	Summary of Changes	Date
Annual Review	<ul style="list-style-type: none"> No clinical policy statement changes. 	1/15/2026
Focused Review	<ul style="list-style-type: none"> Added noncoverage policy statement for intent decoding and pattern recognition add-on modules for upper limb prostheses. 	6/15/2025
Annual Review	<ul style="list-style-type: none"> Removed policy statements for consumable supplies; upper limb sensor and myoelectric controlled prosthetic device with simultaneous multiple degrees of freedom; upper limb prosthetic device using electromyography-based brain computer interface; osseointegrated/osseoanchored lower limb 	3/15/2025

Type of Revision	Summary of Changes	Date
	prosthetic device; and repair/replacement of a prosthetic device. <ul style="list-style-type: none"> • Revised policy statements for iris prostheses; external facial prostheses; and lower limb prostheses. 	
Annual Review	<ul style="list-style-type: none"> • No clinical policy statement changes. 	1/15/2024

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2026 The Cigna Group.