



## Drug Coverage Policy

Effective Date .....4/15/2026

Coverage Policy Number.....IP0405

Policy Title.....Enjaymo

# Hematology – Enjaymo

- Enjaymo® (sutimlimab-jome intravenous infusion – Bioverativ/Sanofi)

### INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer’s particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer’s benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer’s benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see “Coding Information” below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

### Overview

Enjaymo, a classical complement inhibitor, is indicated for the treatment of hemolysis in **cold agglutinin disease** in adults.<sup>1</sup>

### Disease Overview

Cold agglutinin disease is a rare form of autoimmune hemolytic anemia with a prevalence of about 16 per million and an incidence of 1 per million year.<sup>2-4,10</sup> Primary cold agglutinin disease is a B-

cell lymphoproliferative disorder in which autoantibodies are produced against erythrocyte surface antigens. Primary cold agglutinin disease is distinct from secondary disease, termed cold agglutinin syndrome, which can occur with underlying conditions such as malignancy, infection, and autoimmune diseases.<sup>2,3</sup> Diagnosis of cold agglutinin disease is defined by chronic hemolysis, a cold agglutinin titer  $\geq 64$  at 4°C, and typical findings on direct antibody test (DAT), which include strong positivity for complement protein C3d and negativity (or only weak positivity) for immunoglobulin G.<sup>2-4</sup> Secondary causes of cold agglutinin syndrome should be excluded. Importantly, patients without chronic hemolysis or circulatory symptoms do not have cold agglutinin disease, even in the presence of positive DAT.<sup>2</sup> Symptoms include cold-induced circulatory symptoms, which can range from slight acrocyanosis to severe Raynaud phenomena. Anemia is generally considered mild to moderate with a median hemoglobin (Hb) of 8.9 g/dL; however, the lower tertile Hb was 8.0 g/dL and ranged to as low as 4.5 g/dL.<sup>2,4</sup>

### **Clinical Efficacy**

In the CARDINAL trial, patients (n = 24) were required to have a confirmed diagnosis of cold agglutinin disease based on chronic hemolysis, typical DAT findings, and a recent blood transfusion within the prior 6 months.<sup>1,5-7</sup> Patients were also required to have a baseline hemoglobin level < 10 g/dL and total bilirubin above normal. Approximately two-thirds of patients had failed other therapies (e.g., rituximab). The Phase III CADENZA trial (n = 42) also required chronic hemolysis, as well as the DAT and cold agglutinin titer findings described above; however, recent history of blood transfusion was not required.<sup>1,8</sup>

### **Dosing Information**

Dosing is weight-based and is provided for patients weighing  $\geq 39$  kg.<sup>1</sup> For a patient weighing 39 to < 75 kg, the recommended dose is 6,500 mg. For a patient weighing  $\geq 75$  kg, the dose is 7,500 mg. For all patients, the initial dosing frequency is once weekly for 2 weeks, with administration once every 2 weeks (Q2W) thereafter. However, if the interval between doses exceeds 17 days, Enjaymo should be administered once weekly for 2 weeks, returning to Q2W administration thereafter.

### **Guidelines**

An international consensus guideline for autoimmune hemolytic anemias was published in 2020.<sup>9</sup> The guideline was published prior to the approval of Enjaymo and no formal recommendation is made regarding its place in therapy, although positive Phase I data are acknowledged. It is noted that clinical and histological assessment, as well as radiologic examinations as needed, are necessary to rule out cold agglutinin syndrome secondary to malignant disease. Treatment of cold agglutinin syndrome involves supportive care and management of the underlying disease. For treatment of cold agglutinin disease, asymptomatic patients should be managed with watchful waiting. For symptomatic patients (i.e., those with anemia, transfusion, or circulatory symptoms), rituximab is the best-documented first-line treatment and may be given alone or in combination with bendamustine. For second-line treatment, the combination of rituximab plus bendamustine is recommended (if not given in the first-line setting). Alternatively, rituximab monotherapy may be repeated for patients who previously responded for at least 1 year. Rituximab plus fludarabine is an option for fit, elderly patients. There are no evidence-based therapies for the third-line setting.

## **Coverage Policy**

### **POLICY STATEMENT**

Prior Authorization is required for benefit coverage of Enjaymo. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Extended approvals are allowed if the patient continues to meet the Criteria and Dosing. Requests for doses outside of the

established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Enjaymo as well as the monitoring required for adverse events and long-term efficacy, approval requires Enjaymo to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Documentation:** Documentation is required where noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart notes, laboratory tests, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information.

**Enjaymo is considered medically necessary when the following criteria are met:**

**FDA-Approved Indication**

**1. Cold Agglutinin Disease.** Approve for 1 year if the patient meets ALL of the following (A, B, C, D, E, F, G, and H):

**A)** Patient is  $\geq$  18 years of age ; AND

**B)** Patient weighs  $\geq$  39 kg ; AND

**C)** Patient has a history of at least one sign or symptom associated with cold agglutinin disease; AND

Note: Examples include symptomatic anemia (e.g., anemia associated with fatigue, weakness, shortness of breath, heart palpitations, lightheadedness, chest pain), acrocyanosis, Raynaud's syndrome, hemoglobinuria, disabling circulatory symptoms, or a major adverse vascular event (e.g., thrombosis).

**D)** According to the prescriber, the patient has evidence of chronic hemolysis; AND

**E)** Patient meets BOTH of the following diagnostic criteria (i and ii) **[documentation required]**:

i. Direct antibody test strongly positive for C3d and negative or only weakly positive for immunoglobulin G; AND

ii. Cold agglutinin antibody titer is greater than 64 at 4°C (approximately 40°F); AND

**F)** At baseline (prior to the initiation of Enjaymo), patient meets BOTH of the following (i and ii) **[documentation required]**:

i. Hemoglobin  $\leq$  10 g/dL; AND

ii. Total bilirubin above the upper limit of normal, based on the reference range for the reporting laboratory; AND

**G)** According to the prescriber, secondary causes of cold agglutinin syndrome have been excluded; AND

Note: Examples of secondary causes of cold agglutinin syndrome include infection, rheumatologic diseases, and active hematologic malignancies.

**H)** Enjaymo is prescribed by or in consultation with a hematologist.

**Dosing.** Approve ONE of the following dosing regimens (A or B):

**A)** Patient weighs  $\geq$  75 kg: Approve 7,500 mg intravenously not more frequently than once weekly.

**B)** Patient weighs  $<$  75 kg: Approve 6,500 mg intravenously not more frequently than once weekly.

## Conditions Not Covered

Enjaymo for any other use is considered not medically necessary. Criteria will be updated as new published data are available.

## Coding Information

**Note:** 1) This list of codes may not be all-inclusive.  
2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

**Considered Medically Necessary when criteria in the applicable policy statements listed above are met:**

HCPCS Codes	Description
J1302	Injection, sutimlimab-jome, 10 mg

## References

1. Enjaymo® intravenous infusion [prescribing information]. Waltham, MA: Bioverativ/Sanofi; March 2023.
2. Berentsen S, Röth A, Randen U, et al. Cold agglutinin disease: current challenges and future prospects. *J Blood Med.* 2019;10:93-103.
3. Berentsen S. How I treat cold agglutinin disease. *Blood.* 2021;137(10):1295-1303.
4. Swiecicki PL, Hegerova LT, Gertz MA. Cold agglutinin disease. *Blood.* 2013;122(7):1114-1121.
5. Röth A, Barcellini W, D'Sa S, et al. Sutimlimab in cold agglutinin disease. *N Engl J Med.* 2021;384(14):1323-1334.
6. Röth A, Broome CM, Barcellini W, et al. Long-term sutimlimab improves quality of life for patients with cold agglutinin disease: CARDINAL 2-year follow-up. *Blood Adv.* 2023;7(19):5890-5897.
7. Röth A, Barcellini W, D'Sa S, et al. Sustained inhibition of complement C1s with sutimlimab over 2 years in patients with cold agglutinin disease. *Am J Hematol.* 2023;98(8):1246-1253.
8. Röth A, Berentsen S, Barcellini W, D'Sa S, et al. Sutimlimab in patients with cold agglutinin disease: results of the randomized placebo-controlled phase 3 CADENZA trial. *Blood.* 2022;140(9):980-991.
9. Jäger U, Barcellini W, Broome CM, et al. Diagnosis and treatment of autoimmune hemolytic anemia in adults: recommendations from the First International Consensus Meeting. *Blood Rev.* 2020 May;41:100648.
10. Barcellini W, Fattizzo B. The evolving management algorithm for a patient with newly diagnosed cold agglutinin disease. *Expert Rev Hematol.* 2024;17(7):287-294.

## Revision Details

Type of Revision	Summary of Changes	Date
Annual Revision	No criteria changes.	5/1/2024
Annual Revision	<b>Added</b> definition of "documentation".	5/1/2025

	Updated "documentation" phrasing throughout coverage policy.	
Annual Revision	<p><b>Added</b> the following note to provide examples of signs and symptoms associated with cold agglutinin disease. "Note: Examples include symptomatic anemia (e.g., anemia associated with fatigue, weakness, shortness of breath, heart palpitations, lightheadedness, chest pain), acrocyanosis, Raynaud's syndrome, hemoglobinuria, disabling circulatory symptoms, or a major adverse vascular event (e.g., thrombosis)."</p> <p><b>Added</b> the following note to provide examples of secondary causes of cold agglutinin syndrome. "Note: Examples of secondary causes of cold agglutinin syndrome include infection, rheumatologic diseases, and active hematologic malignancies."</p> <p><b>Dosing:</b> Updated dosing from "Weight of 39 kg to less than 75 kg" to "Patient weighs &lt; 75 kg"</p> <p><b>Conditions Not Covered:</b> Removed Paroxysmal Cold Hemoglobinuria.</p>	04/15/2026

The policy effective date is in force until updated or retired.

---

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2026 The Cigna Group.