



Drug Coverage Policy

Effective Date..... 04/01/2026

Coverage Policy Number IP0433

Policy Title.....Symdeko

Cystic Fibrosis Transmembrane Conductance Regulator – Symdeko

- Symdeko® (tezacaftor/ivacaftor and ivacaftor tablets - Vertex)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

OVERVIEW

Symdeko, a cystic fibrosis transmembrane conductance regulator (CFTR), is indicated for the treatment of **cystic fibrosis** (CF) in patients \geq 6 years of age who are homozygous for the

F508del mutation or who have at least one mutation in the *CFTR* gene that is responsive to tezacaftor/ivacaftor based on *in vitro* data and/or clinical evidence.¹

If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a *CFTR* mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use. Table 1 lists the responsive *CFTR* mutations based on: 1) a clinical forced expiratory volume in 1 second (FEV₁) response and/or 2) *in vitro* data in Fischer rat thyroid cells, indicating that tezacaftor/ivacaftor increases chloride transport to ≥ 10% of untreated normal over baseline. *CFTR* gene mutations that are not responsive to Kalydeco® (ivacaftor granule or tablet) alone are not expected to respond to Symdeko except for *F508del* homozygotes.

Table 1. List of CFTR Gene Mutations that Produce CFTR Protein and are Responsive to Symdeko.¹

<i>E56K</i>	<i>E193K</i>	<i>S945L</i>	<i>F1074L</i>
<i>P67L</i>	<i>L206W</i>	<i>S977F</i>	<i>D1152H</i>
<i>R74W</i>	<i>R347H</i>	<i>F1052V</i>	<i>D1270N</i>
<i>D110E</i>	<i>R352Q</i>	<i>E831X</i>	<i>2789+5G→A</i>
<i>D110H</i>	<i>A455E</i>	<i>K1060T</i>	<i>3272-26A→G</i>
<i>R117C</i>	<i>D579G</i>	<i>A1067T</i>	<i>3849 + 10kbC→T</i>
<i>F508del*</i>	<i>711+3A→G</i>	<i>R1070W</i>	<i>G622D</i>
<i>A120T</i>	<i>E60K</i>	<i>F1016S</i>	<i>G970D</i>
<i>A234D</i>	<i>E92K</i>	<i>F1099L</i>	<i>G1069R</i>
<i>A349V</i>	<i>E116K</i>	<i>G126D</i>	<i>G1244E</i>
<i>A554E</i>	<i>E403D</i>	<i>G178E</i>	<i>G1249R</i>
<i>A1006E</i>	<i>E558V</i>	<i>G178R</i>	<i>G1349D</i>
<i>D192G</i>	<i>E822K</i>	<i>G194R</i>	<i>H939R</i>
<i>D443Y</i>	<i>F191V</i>	<i>G194V</i>	<i>H1054D</i>
<i>D443Y;G57A; R668C</i>	<i>F311del</i>	<i>G314E</i>	<i>H1375P</i>
<i>D614G</i>	<i>F311L</i>	<i>G551D</i>	<i>I148T</i>
<i>D836Y</i>	<i>F508C</i>	<i>G551S</i>	<i>I175V</i>
<i>D924N</i>	<i>F508C;S1251N</i>	<i>G576A</i>	<i>I336K</i>
<i>D979V</i>	<i>F575Y</i>	<i>G576A;R668C</i>	<i>I601F</i>
<i>I618T</i>	<i>L346P</i>	<i>M952T</i>	<i>R74Q</i>
<i>I807M</i>	<i>L967S</i>	<i>P5L</i>	<i>R74W;D1270N</i>
<i>I980K</i>	<i>L997F</i>	<i>P205S</i>	<i>R74W;V201M</i>
<i>I1027T</i>	<i>L1324P</i>	<i>Q98R</i>	<i>R74W;V201M;D1270N</i>
<i>I1139V</i>	<i>L1335P</i>	<i>Q237E</i>	<i>R75Q</i>
<i>I1269N</i>	<i>L1480P</i>	<i>Q237H</i>	<i>R117G</i>
<i>I1366N</i>	<i>M152V</i>	<i>Q359R</i>	<i>R117H</i>
<i>L15P</i>	<i>M265R</i>	<i>Q1291R</i>	<i>R117L</i>
<i>L320V</i>	<i>M952I</i>	<i>R31L</i>	<i>R117P</i>
<i>R170H</i>	<i>R1066H</i>	<i>S1251N</i>	<i>W1282R</i>
<i>R258G</i>	<i>R1070Q</i>	<i>S1255P</i>	<i>Y109N</i>
<i>R334L</i>	<i>R1162L</i>	<i>T338I</i>	<i>Y161S</i>
<i>R334Q</i>	<i>R1283M</i>	<i>T1036N</i>	<i>Y1014C</i>
<i>R347L</i>	<i>R1283S</i>	<i>T1053I</i>	<i>Y1032C</i>
<i>R347P</i>	<i>S549N</i>	<i>V201M</i>	<i>R792G</i>
<i>R352W</i>	<i>S549R</i>	<i>V232D</i>	<i>R933G</i>
<i>R553Q</i>	<i>S589N</i>	<i>V562I</i>	<i>S1159F</i>

<i>R668C</i>	<i>S737F</i>	<i>V754M</i>	<i>S1159P</i>
<i>R751L</i>	<i>S912L</i>	<i>V1153E</i>	<i>V1240G</i>
<i>V1293G</i>	<i>546insCTA</i>		

CFTR – Cystic fibrosis transmembrane conductance regulator; * A patient must have two copies of the F508del mutation or at least one copy of a responsive mutation presented in Table 1 to be indicated.

Guidelines

The Standards of Care for CFTR variant-specific therapy for people with CF, from the European Cystic Fibrosis Society (2023) do not reflect the currently approved age indications for Kalydeco [≥ 1 months of age], Orkambi® [lumacaftor/ivacaftor tablets and oral granules] (≥ 1 year of age), or Trikafta® (elexacaftor/tezacaftor/ivacaftor; ivacaftor co-packaged tablets and granules) [≥ 2 years of age].² In general, Trikafta is recommended over other agents where indications overlap. The Standards recommend Trikafta in patients ≥ 6 years of age with CF who are homozygous or heterozygous for *F508del*. In patients with one or more responsive non-*F508del* variant, Kalydeco, Symdeko, or Trikafta are recommended. Kalydeco is recommended in patients ≥ 4 months of age with eligible *CFTR* gene variants. Orkambi is recommended for patients 2 to 5 years of age who are homozygous for *F508del*. Of note, the Standards state that after diagnosis, repeat sweat testing provides evidence of treatment effect on CFTR activity, but does not predict clinical response. The European Cystic Fibrosis Society Standards for establishing and maintaining health (2024) note that people with CF with eligible *CFTR* gene variants should be offered CFTR modulator therapy.⁵

According to the CF Foundation (2017), CF is diagnosed when an individual has both a clinical presentation of CF and evidence of CFTR dysfunction.^{3,4} Clinical presentation of CF includes a positive newborn screening, signs and/or symptoms of CF, and/or family history of CF. To establish a diagnosis of CF, sweat chloride tests should be considered first, then *CFTR* genetic analysis (*CFTR* genotype), and then CFTR physiologic tests (nasal potential difference [NPD] or intestinal current measurement [ICM]). However, tests of CFTR function are not always done in this order. All individuals diagnosed with CF should have a sweat chloride test and *CFTR* genetic analysis performed.

In a patient with a sweat chloride test ≥ 60 mmol/L, CF diagnosis is established and in patients with a sweat chloride test < 30 mmol/L, a diagnosis of CF is unlikely.^{3,4} Rarely, patients with a sweat chloride < 30 mmol/L may be considered to have CF if alternatives are excluded and other confirmatory tests (genetic and physiologic testing) support a CF diagnosis. In patients with a sweat chloride test of ≥ 30 to < 60 mmol/L, CFTR genetic analysis is undertaken. If the genetic analysis identifies two CF-causing *CFTR* mutations, CF is diagnosed; if no *CFTR* mutations are identified, a diagnosis of CF is unlikely. In patients with a *CFTR* genotype that is undefined or of varying clinical consequence, full gene *CFTR* sequencing (if not already performed) or CFTR physiologic testing is performed (NPD or ICM). If only one *CFTR* variant is identified on limited analysis, full gene *CFTR* sequencing should be performed. CF is possible if both alleles possess CF-causing, undefined, or mutation of varying clinical consequence mutations; CF is unlikely if only no CF-causing mutations are found. If results of the NPD or ICM show CFTR dysfunction, CF is diagnosed; when testing is unavailable or equivocal, the diagnosis of CF is not resolved, and when results of the physiologic testing show CFTR function is preserved, a diagnosis of CF is considered unlikely. It is recommended that patients with challenging diagnoses be evaluated at an accredited CF Foundation Care Center.

Coverage Policy

POLICY STATEMENT

Prior Authorization is required for benefit coverage of Symdeko. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Symdeko as well as the monitoring required for adverse events and efficacy, approval requires Symdeko to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Documentation: Documentation is required where noted in the criteria as **[documentation required]**. Documentation may include, but not limited to, chart notes, laboratory tests, medical test results, claims records, and/or other information. All documentation must include patient-specific identifying information.

Symdeko is considered medically necessary when the following criteria are met:

- 1. Cystic Fibrosis.** Approve for 1 year if the patient meets the following (A, B, C D, E, and F):
 - A)** Patient is ≥ 6 years of age; AND
 - B)** Patient meets ONE of the following (i or ii):
 - i.** Patient has at least ONE of the following variants in the cystic fibrosis transmembrane conductance regulator gene that is considered to be a pathogenic or likely pathogenic variant: *E56K, P67L, R74W, D110E, D110H, R117C, E193K, L206W, R347H, R352Q, A455E, D579G, 711+3A \rightarrow G, S945L, S977F, F1052V, E831X, K1060T, A1067T, R1070W, F1074L, D1152H, D1270N, 2789+5G \rightarrow A, 3272-26A \rightarrow G, 3849 + 10kbc \rightarrow T, 546insCTA, A120T, A234D, A349V, A554E, A1006E, D192G, D443Y, D443Y;G57A;R668C, D614G, D836Y, D924N, D979V, I618T, I807M, I980K, I1027T, I1139V, I1269N, I1366N, L15P, L320V, R170H, R258G, R334L, R334Q, R347L, R347P, R352W, R553Q, R668C, R751L, V1293G, E60K, E92K, E116K, E403D, E558V, E822K, F191V, F311del, F311L, F508C, F508C;S1251N, F575Y, L346P, L967S, L997F, L1324P, L1335P, L1480P, M152V, M265R, M952I, R1066H, R1070Q, R1162L, R1283M, R1283S, S549N, S549R, S589N, S737F, S912L, F1016S, F1099L, G126D, G178E, G178R, G194R, G194V, G314E, G551D, G551S, G576A, G576A;R668C, M952T, P5L, P205S, Q98R, Q237E, Q237H, Q359R, Q1291R, R31L, S1251N, S1255P, T338I, T1036N, T1053I, V201M, V232D, V562I, V754M, V1153E, G622D, G970D, G1069R, G1244E, G1249R, G1349D, H939R, H1054D, H1375P, I148T, I175V, I336K, I601F, R74Q, R74W;D1270N, R74W;V201M, R74W;V201M;D1270N, R75Q, R117G, R117H, R117L, R117P, W1282R, Y109N, Y161S, Y1014C, Y1032C, R792G, R933G, S1159F, S1159P, or V1240G **[documentation required]**; OR*
 - ii.** Patient has TWO copies of the *F508del* variant; AND
 - C)** Patient meets at least ONE of the following (i, ii, or iii):
 - i.** Positive cystic fibrosis newborn screening test; OR
 - ii.** Family history of cystic fibrosis; OR
 - iii.** Clinical presentation consistent with signs and symptoms of cystic fibrosis; AND
Note: Examples of clinical presentation of cystic fibrosis include but are not limited to meconium ileus, sino-pulmonary symptoms (e.g., persistent cough, wheezing, pulmonary function tests consistent with obstructive airway disease, excess sputum production), bronchiectasis, sinusitis, failure to thrive, pancreatic insufficiency.
 - D)** Patient has evidence of abnormal cystic fibrosis transmembrane conductance regulator function as demonstrated by at least ONE of the following (i, ii, or iii):
 - i.** Elevated sweat chloride test; OR
 - ii.** Two cystic fibrosis-causing cystic fibrosis transmembrane conductance regulator variants; OR
 - iii.** Abnormal nasal potential difference; AND
 - E)** The medication is prescribed by or in consultation with a pulmonologist or a physician who specializes in the treatment of cystic fibrosis.
 - F)** Preferred product criteria is met for the products listed in the below table(s)

Employer Plans:

Product	Criteria
Symdeko (tezacaftor/ivacaftor and ivacaftor tablets)	Total Savings Drug List Plans: Patient meets ONE of the following (1, 2, <u>or</u> 3): 1. Patient has tried, and according to the prescriber has experienced inadequate efficacy OR a significant intolerance with Trikafta (tablets or oral granules) [may require prior authorization] 2. Patient has at least one variant in the cystic fibrosis transmembrane regulator gene that is considered to be a pathogenic or likely pathogenic variant that is not covered by Trikafta (tablets or granules) [may require prior authorization] 3. Patient has already been started on therapy with Symdeko

Symdeko for any other use is considered not medically necessary, including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

- 1. Cystic Fibrosis, Patients with Unknown Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) Gene Mutation.** If the patient has an unknown genotype, an FDA-cleared CF mutation test should be used to detect the presence of the cystic fibrosis transmembrane regulator variant followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.¹
- 2. Combination Therapy with Other Cystic Fibrosis Transmembrane Conductance Regulator Modulator(s).** Symdeko contains ivacaftor, the active agent in Kalydeco® (tablets and oral granules) and part of Orkambi® (lumacaftor/ivacaftor tablets and oral granules) and Trikafta® (elexacaftor/tezacaftor/ivacaftor; ivacaftor co-packaged tablets and granules). Symdeko also contains tezacaftor, part of Trikafta.
Note: Examples of other cystic fibrosis transmembrane conductance regulator modulators are: Alyftrek™ (vanzacaftor/tezacaftor/deutivacaftor tablets), Kalydeco (ivacaftor tablets and oral granules), Orkambi (lumacaftor/ivacaftor tablets and oral granules), Trikafta (elexacaftor/tezacaftor/ivacaftor; ivacaftor co-packaged tablets and granules).
- 3. Infertility.** Symdeko is indicated for the treatment of cystic fibrosis in a patient ≥ 6 years of age who is homozygous for the *F508del* variant or who has at least one variant in the cystic fibrosis transmembrane conductance regulator gene that is responsive to tezacaftor/ivacaftor based on *in vitro* data and/or clinical evidence.¹
Note: A patient with a diagnosis of cystic fibrosis should be reviewed using criteria for the FDA-approved indication, above.

References

1. Symdeko® tablets [prescribing information]. Cambridge, MA: Vertex; September 2025.
2. Southern KW, Addy C, Bell SC, et al. Standards for the care of people with cystic fibrosis; establishing and maintaining health. *J Cyst Fibros.* 2024;21-28.
3. Farrell PM, White TB, Ren CL, et al. Diagnosis of cystic fibrosis: consensus guidelines from the cystic fibrosis foundation. *J Pediatr.* 2017;181S:S4-S15.

4. Farrell PM, White TB, Howenstine MS, et al. Diagnosis of cystic fibrosis in screened populations. *J Pediatr.* 2017;181S:S33-S44.
5. Southern KW, Addy C, Bell SC, et al. Standards for the care of people with cystic fibrosis; establishing and maintaining health. *J Cyst Fibros.* 2024;21-28.

Revision Details

Type of Revision	Summary of Changes	Date
Annual Revision	<p>Updated coverage policy title from Tezacaftor/ Ivacaftor to Cystic Fibrosis – Symdeko.</p> <p>Added preferred product criterion to allow coverage Symdeko if there is at least one of the following mutations in the CFTR gene: 711+3A > G, E831X, 2789+5G > A, 3272-26A > G, OR 3849 + 10kbC > T</p> <p>Specified Preferred Product Table applies to Total Savings Drug List Plans.</p> <p>Infertility: Added this indication to conditions not recommended for approval.</p> <p>CFTR-related disorder: Removed indication from <i>Conditions Not Covered</i>.</p> <p>CFTR-related metabolic syndrome, CF Screen Positive, Inconclusive Diagnosis (CRMS/CF SPID): Removed indications from <i>Conditions Not Covered</i>.</p>	08/01/2024
Selected Revision	<p>The Policy title was changed to Cystic Fibrosis Transmembrane Conductance Regulator – Symdeko. Previously, Cystic Fibrosis – Symdeko.</p> <p>Added “Documentation: Documentation is required where noted in the criteria. Documentation may include, but not limited to, chart notes, laboratory tests, medical test results, claims records, and/or other information.”</p> <p>Cystic Fibrosis: Updated criteria from “Patient meets ONE of the following (i <u>or</u> ii):” to “Documentation is provided that the patient meets ONE of the following (i <u>or</u> ii):”</p> <p>Cystic Fibrosis, Patient with Unknown Cystic Fibrosis Transmembrane Conductance Regulator Gene Mutation. “Conductance” was added to the verbiage for this condition not covered.</p>	04/01/2025

	<p>Combination Therapy with Other Cystic Fibrosis Transmembrane Conductance Regulator Modulator(s). This condition not covered was modified to refer to the class of cystic fibrosis transmembrane conductance regulator modulator(s). Previously individual agents were listed. A Note was added to list examples of the cystic fibrosis transmembrane conductance regulators.</p> <p>Preferred Product Table: Updated from "Failure, contraindication, or intolerance to elexacaftor/tezacaftor/ ivacaftor (Trikafta™)" to "Patient has tried and according to the prescriber has experienced inadequate efficacy OR a significant intolerance with Trikafta(tablets or oral granules) [may require prior authorization]."Added "Patient has at least one mutation in the cystic fibrosis transmembrane regulator gene that is considered to be a pathogenic or likely pathogenic variant that is not covered by Trikafta (tablets or granules) [may require prior authorization]." Removed "At least ONE of the following mutations in the cystic fibrosis transmembrane regulator (CFTR) gene: 711+3A > G, E831X, 2789+5G > A, 3272-26A > G, OR 3849 + 10kbC > T." Updated from "Currently receiving Symdeko" to "Patient has already been started on therapy with Symdeko."</p>	
Annual Revision	<p>Updated the documentation requirements to current standards.</p> <p>Cystic Fibrosis. The term "mutation" was replaced by "variant" for the following requirements: The patient has at least ONE of the following variants in the cystic fibrosis transmembrane conductance regulator gene that is considered pathogenic or likely pathogenic; the patient has TWO copies of the <i>F508del</i> variant; and the patient has evidence of abnormal cystic fibrosis transmembrane conductance regulator function as demonstrated by two cystic fibrosis-causing cystic fibrosis transmembrane conductance regulator variants.</p> <p>Conditions Not Recommended for Approval Cystic Fibrosis, Patient with Unknown Cystic Fibrosis Transmembrane Conductance Regulator Gene Variant. The term "mutation"</p>	04/01/2026

	was replaced by "variant" in this condition not recommended for approval.	
--	---	--

The policy effective date is in force until updated or retired.

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2026 The Cigna Group.