



## Drug Coverage Policy

Effective Date .....04/30/2026  
Coverage Policy Number.....IP0739  
Policy Title.....Weight Loss –  
Glucagon-Like Peptide-1 Agonists BMI  
≥ 35

# Weight Loss – Glucagon-Like Peptide-1 Agonists BMI ≥ 35

- Foundayo™ (orforglipron tablets – Eli Lilly)
- Saxenda® (liraglutide subcutaneous injection – Novo Nordisk, generic)
- Wegovy® (semaglutide tablet and subcutaneous injection – Novo Nordisk)
- Zepbound™ (tirzepatide subcutaneous injection – Eli Lilly)

---

### **INSTRUCTIONS FOR USE**

*The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used*

as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

---

Weight loss medications are specifically excluded under many benefit plans [both Employer Groups and Individual and Family Plans]. Please refer to the applicable benefit plan document to determine benefit availability and the terms and conditions of coverage.

### Overview

Liraglutide (Saxenda, generic), Foundayo, Wegovy, and Zepbound are glucagon-like peptide-1 (GLP-1) receptor agonists; Zepbound is also a glucose-dependent insulinotropic polypeptide (GIP) receptor agonist.<sup>1,2,4,25</sup>

**Liraglutide, Foundayo, Wegovy injection, Wegovy tablet, and Zepbound** are indicated in combination with a reduced-calorie diet and increased physical activity:<sup>1,2,4,25</sup>

- To **reduce excess body weight and maintain weight reduction long term** in:
  - **Liraglutide, Foundayo, Wegovy injection, Wegovy tablet, and Zepbound:** Adults with overweight in the presence of at least one weight-related comorbid condition.<sup>1,2,4,6,25</sup>
  - **Liraglutide, Foundayo, Wegovy injection, Wegovy tablet, and Zepbound:** Adults with obesity.<sup>1,4,25</sup>
  - **Liraglutide:** Pediatric patients  $\geq 12$  years of age and  $\geq 60$  kg with obesity.<sup>2</sup>
  - **Wegovy injection:** Pediatric patients  $\geq 12$  years of age with obesity.<sup>1,7</sup>

**Wegovy injection and Wegovy tablet** are indicated in combination with a reduced-calorie diet and increased physical activity:<sup>1</sup>

- To **reduce the risk of major adverse cardiovascular (CV) events (MACE)** [CV death, non-fatal myocardial infarction {MI}, or non-fatal stroke] in adults with established CV disease and either **obesity or overweight**.<sup>1,5</sup>

**Wegovy injection** is indicated in combination with a reduced-calorie diet and increased physical activity:<sup>1</sup>

- For the treatment of **non-cirrhotic metabolic dysfunction-associated steatohepatitis (MASH)**, formerly known as non-alcoholic steatohepatitis (NASH), with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in adults.<sup>1,19,20</sup>

**Zepbound** is indicated in combination with a reduced-calorie diet and increased physical activity:<sup>4</sup>

- To treat **moderate to severe obstructive sleep apnea (OSA)** in adults with **obesity**.

### Dosing

In the prescribing information for Foundayo, the recommended starting dose is 0.8 mg orally once daily (QD).<sup>25</sup> After  $\geq 30$  days on the 0.8 mg dose, increase the dose to 2.5 mg QD. After  $\geq 30$  days on the 2.5 mg dose, increase the dose to 5.5 mg QD. The dose may be increased to the next dose level (9 mg, 14.5 mg, or 17.2 mg QD) after  $\geq 30$  days on the current dose based on treatment response and tolerability. The maximum dose is 17.2 mg QD (reached after  $\geq 180$  days).

In the prescribing information for Wegovy injection, a recommended dose escalation schedule of 16 weeks is outlined (the 2.4 mg dose would be reached at the start of Week 17).<sup>1</sup> If a patient does not tolerate a dose during dose escalation, consider delaying dose escalation for 4 weeks.

In the prescribing information for Wegovy tablet, a recommended dose escalation schedule of 90 days is outlined (the 25 mg dose would be reached at the start of Day 91).<sup>1</sup> If a patient does not tolerate a dose during dose escalation, consider delaying dose escalation. For CV risk reduction and weight reduction, the maintenance dose of Wegovy tablet is 25 mg orally QD. If the patient does not tolerate the 25 mg QD maintenance dosage, consider switching to Wegovy subcutaneous (SC) injection 1.7 mg once weekly (QW). If additional weight reduction is needed in patients with type 2 diabetes mellitus treated with Wegovy 25 mg tablet, consider switching to Wegovy 1.7 mg SC injection QW and follow the recommended dosage escalation for Wegovy SC injection.

Adults taking Wegovy 2.4 mg SC injection for CV risk reduction or weight reduction may switch to Wegovy 25 mg tablets.<sup>1</sup> One week after discontinuing Wegovy 2.4 mg SC injection, initiate 25 mg of Wegovy tablets orally QD. A patient may switch from Wegovy 25 mg tablets to Wegovy SC injection. The day after discontinuing Wegovy 25 mg tablets QD, initiate Wegovy 2.4 mg SC injection QW.

In the prescribing information for liraglutide, a recommended dose escalation schedule of 4 weeks is outlined.<sup>2</sup> If a patient does not tolerate an increased dose during dose escalation, consider delaying dose escalation for approximately one additional week. For adults, the recommended maintenance dose of liraglutide is 3 mg SC QD; discontinue liraglutide if the patient cannot tolerate the 3 mg dose. Additionally, for adults, the prescribing information states to evaluate the change in body weight 16 weeks after initiating liraglutide and discontinue liraglutide if the patient has not lost  $\geq 4\%$  of baseline body weight, since it is unlikely the patient will achieve and sustain clinically meaningful weight loss with continued treatment. For pediatric patients, the recommended maintenance dose is 3 mg SC QD; patients who do not tolerate 3 mg QD may have the maintenance dose reduced to 2.4 mg QD. Discontinue liraglutide if the patient cannot tolerate the 2.4 mg dose. If a pediatric patient does not tolerate an increased dose during dose escalation, the dose may be lowered to the prior dose level; dose escalation may take up to 8 weeks. Evaluate the change in body mass index (BMI) after 12 weeks on the maintenance dose in pediatric patients; if the patient has not had a reduction in BMI of  $\geq 1\%$  from baseline, discontinue liraglutide as it is unlikely the patient will achieve and sustain clinically meaningful weight loss with continued treatment.

In the prescribing information for Zepbound, the recommended starting dose is 2.5 mg SC QW.<sup>4</sup> The 2.5 mg dose is for treatment initiation and is not intended for chronic weight management. After 4 weeks, the dose can be increased to 5 mg QW. The dose can then be increased in 2.5 mg increments, after at least 4 weeks on the current dose. The recommended maintenance doses for weight reduction and long-term maintenance are 5 mg, 10 mg, or 15 mg QW. The recommended maintenance dose in OSA is 10 mg or 15 mg QW. The treatment response and tolerability should be considered when selecting the maintenance dose. If a patient does not tolerate a maintenance dose, consider a lower maintenance dose. The maximum dose is 15 mg QW. The 5 mg, 10 mg, and 15 mg maintenance doses are reached after Week 4, Week 12, and Week 20, respectively.

None of the GLP-1 or GLP-1/GIP agonists are recommended for coadministration with other GLP-1 or GLP-1/GIP agonists.<sup>1,2,4,25</sup>

## **Clinical Efficacy**

### *Secondary Prevention of MACE*

SELECT was a randomized, double-blind, placebo-controlled, event-driven study that assessed Wegovy injection vs. placebo, when added to standard of care, for the secondary prevention of CV

events in adults  $\geq 45$  years of age with BMI  $\geq 27$  kg/m<sup>2</sup> and established CV disease without diabetes (n = 17,604).<sup>5</sup> Established CV disease was defined as one of the following: prior MI, prior stroke (ischemic or hemorrhagic), and/or symptomatic peripheral arterial disease (as evidenced by intermittent claudication with ankle-brachial index  $< 0.85$ , peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease). The primary efficacy endpoint was a composite of death from CV causes, non-fatal MI, or non-fatal stroke.

**Results.** Patients were followed for a mean of 39.8 months.<sup>5</sup> The trial achieved its primary endpoint, demonstrating a statistically significant and superior reduction in MACE for Wegovy injection vs. placebo. A primary endpoint event occurred in 6.5% vs. 8.0% of patients in the Wegovy injection vs. placebo groups, respectively (hazard ratio 0.80; 95% confidence interval: 0.72, 0.90; P < 0.001).

#### *MASH*

The ESSENCE trial (Part 1 n = 800), a two-part, ongoing, Phase III, multicenter, double-blind, parallel-group trial randomized adults with MASH and stage F2 to F3 fibrosis to Wegovy injection or placebo, both in addition to standard of care (optimization of treatment for type 2 diabetes, dyslipidemia, and CV risk management).<sup>19,20</sup> Results from Part 1 have been published. Eligible patients were  $\geq 18$  years of age with histological presence of steatohepatitis with stage F2 to F3 fibrosis from a baseline liver biopsy. Patients with an average alcohol consumption of  $\geq 20$  grams/day for women or  $\geq 30$  grams/day for men or alcohol dependence were excluded. Rezdifra™ (resmetirom tablets) was not approved at the time the trial commenced; therefore, no patients were taking Rezdifra in Part 1 of this trial. Concomitant use of any other GLP-1 or GLP-1/GIP agonist was not allowed. The two primary histologic endpoints were: 1) Resolution of steatohepatitis and no worsening of liver fibrosis; and 2) Improvement in liver fibrosis and no worsening of steatohepatitis. In Part 2 of the trial, the primary endpoint will be cirrhosis-free survival at Week 240 (ongoing). Overall, 56% of patients had type 2 diabetes. The mean BMI was 34.6 kg/m<sup>2</sup>. Most patients fulfilled four (27.8%) or five (43.3%) of five metabolic dysfunction-associated metabolic liver disease (MASLD) cardiometabolic criteria (i.e., BMI  $\geq 25$  kg/m<sup>2</sup> [ $\geq 23$  kg/m<sup>2</sup> Asia] or waist circumference  $> 94$  cm [male] or  $> 80$  cm [female] or ethnicity adjusted equivalent; fasting serum glucose  $\geq 100$  mg/dL or 2-hour post-prandial glucose  $\geq 140$  mg/dL or type 2 diabetes or treatment for type 2 diabetes; blood pressure  $\geq 130/85$  mmHg or specific antihypertensive drug treatment; plasma high-density lipoprotein cholesterol  $\leq 40$  mg/dL [male] and  $\leq 50$  mg/dL [female] or lipid-lowering treatment). Most patients had stage F3 fibrosis (68.8%); 31.3% of patients had stage F2 fibrosis.

**Results.** At the interim analysis (the first 800 patients enrolled in the trial), the between-group differences for both primary endpoints were significant for Wegovy injection vs. placebo.<sup>20</sup> Wegovy injection demonstrated a significant improvement in liver fibrosis with no worsening of steatohepatitis, as well as resolution of steatohepatitis with no worsening of liver fibrosis. Confirmatory secondary endpoints also generally favored Wegovy injection (e.g., resolution of steatohepatitis with improvement in liver fibrosis, weight change). Part 2 of the trial is ongoing and expected to read out in 2029.

#### *OSA*

The SURMOUNT-OSA (n = 469) trials were two 52-week, Phase III, multicenter, double-blind, randomized trials that evaluated the efficacy and safety of maximally tolerated Zepbound (10 mg or 15 mg QW) in adults with obesity (without diabetes) and moderate to severe OSA.<sup>9</sup> Two patient populations were included. In Trial 1, patients were unable or unwilling to use positive airway pressure (PAP) therapy, and in Trial 2, patients had been using PAP therapy for  $\geq 3$  months at the time of screening and planned to continue PAP therapy during the trial. All patients had a diagnosis of moderate to severe OSA with an apnea-hypopnea index (AHI)  $\geq 15$  events/hour as diagnosed with polysomnography, home sleep apnea test, or other methods that met local

guidelines prior to Visit 1. Patients had a BMI of  $\geq 30$  kg/m<sup>2</sup> ( $\geq 27$  kg/m<sup>2</sup> in Japan) despite the history of at least one self-reported unsuccessful dietary effort to lose weight. Patients with a diagnosis of central or mixed sleep apnea with the percentage of mixed or central apneas/hypopneas  $\geq 50\%$ , and Cheyne Stokes respiration were excluded. In Trial 1, the mean BMI was 39.1 kg/m<sup>2</sup> and the mean AHI was 51.5 events/hour. Most patients had severe OSA (63%). In Trial 2, the mean BMI was 38.7 kg/m<sup>2</sup> and the mean AHI was 49.5 events/hour. Most patients had severe OSA (68%). The primary endpoint was the superiority of Zepbound vs. placebo for the change in the AHI from baseline.<sup>9</sup> Several key secondary endpoints were assessed.

**Results.** In both trials, Zepbound was superior to placebo for the primary endpoint. In Trial 1, the change in AHI at Week 52 with Zepbound was superior to placebo (-25.3 events/hour vs. -5.3 events/hour, respectively; estimated treatment difference of -20.0 events/hour;  $P < 0.001$ ).<sup>9</sup> In Trial 2, the change in AHI at Week 52 with Zepbound was superior to placebo (-29.3 events/hour vs. -5.5 events/hour, respectively; estimated treatment difference -23.8 events/hour;  $P < 0.001$ ). Additionally, patients in both trials who received Zepbound had significant reductions in sleep apnea-specific hypoxic burden. The proportion of patients with a reduction in the AHI of  $\geq 50\%$  at Week 52 and the proportion of patients with an AHI of  $< 5$  events/hour or an AHI of 5 to 14 events/hour and an ESS of  $\leq 10$  at Week 52 also favored Zepbound. Patients receiving Zepbound in both trials had significant reductions in body weight.

## Guidelines

Foundayo and Wegovy tablet have not been addressed in guidelines.

### *Weight Management*

#### Adult

The American Academy of Clinical Endocrinology (AACE) Consensus Statement: Algorithm for the Evaluation and Treatment of Adults with Obesity/Adiposity-Based Chronic Disease (ABCD) [2025 update] places an emphasis on a complication-centric, person-centered care model.<sup>23</sup> BMI is appropriate to screen for ABCD/obesity and is used to classify individuals into categories of overweight (BMI  $\geq 25.0$  to  $\leq 29.9$  kg/m<sup>2</sup>), Class I obesity (BMI  $\geq 30.0$  to  $\leq 34.9$  kg/m<sup>2</sup>), Class II obesity (BMI  $\geq 35.0$  to  $\leq 39.9$  kg/m<sup>2</sup>), or Class III obesity (BMI  $\geq 40$  kg/m<sup>2</sup>). Pharmacotherapy, in adjunct to lifestyle modification, is indicated to prevent, improve, or reverse obesity-related complications and diseases; not solely to reduce BMI. The choice of pharmacotherapy is based on obesity-related comorbidities. The degree of weight reduction with a given medication can serve as a guide toward improvement of various comorbidities. Response to pharmacologic therapy to therapy should be assessed after 3 months on a therapeutic dose. If treatment has not resulted in  $\geq 5\%$  weight loss, longer-term efficacy will not likely be sufficient; a change in therapeutic approach is recommended. Individuals with weight reduction  $\geq 5\%$  should continue with their current treatment. Patients who achieve  $\geq 15\%$  weight loss (noted to be the percent weight loss observed on average with Wegovy injection and Zepbound) will have achieved a response to therapy that predictably prevents or improves a wide range of obesity-related comorbidities.

The American Diabetes Association Professional Practice Committee for Obesity (2026) Standards of Care in Overweight and Obesity prioritize obesity pharmacotherapy that is most likely to achieve and maintain intended treatment goals.<sup>24</sup> The treatment goal should be individualized based on the severity of obesity, obesity-related complications and diseases, and individual needs. A sustained weight reduction of  $\geq 5\%$  from baseline may achieve some clinically meaningful health benefits, while a sustained weight reduction of  $\geq 10\%$  is recommended to manage many obesity-related diseases or complications. In some cases,  $\geq 15\%$  weight reduction may be indicated to achieve greater therapeutic benefit. Zepbound and Wegovy injection (listed in order of relative benefit) are recognized as agents with high efficacy ( $> 10\%$  weight loss). Medications should be continued after reaching treatment goals to maintain health benefits. When there is an inadequate response to therapy, dose escalation to the maximum dose is recommended. If the

maximum dose has been reached and the response is inadequate, switching to an alternative medication, adding intensive lifestyle therapy, combining medications, or referring for metabolic surgery is recommended. Patients who achieve early weight reduction (usually  $\geq 5\%$  after 3 months) should continue medication long term.

### Pediatric

Guidelines from the American Academy of Pediatrics on evaluation and treatment of children and adolescents with obesity (2023) note that pediatricians and other primary healthcare providers should offer adolescents  $\geq 12$  years of age with obesity (BMI  $\geq 95^{\text{th}}$  percentile) weight loss pharmacotherapy, according to medication indications, risks, and benefits, as an adjunct to health behavior and lifestyle treatment.<sup>7</sup>

### *MASH*

The American Association for the Study of Liver Diseases (AASLD) Practice Guidance on the Clinical Management of non-alcoholic fatty liver disease (2023) was updated in October 2024 to address the approval of Rezdiffra and in November 2025 to address approval of Wegovy for MASH.<sup>21,22</sup> Best practices in the management of MASH include comprehensive lifestyle modification (nutrition, exercise, behavioral modification), and optimal control of comorbid metabolic conditions.<sup>21</sup> MASH can only be definitively diagnosed by histologic exam (biopsy); however, in practice, patient selection is based on evidence of steatosis and fibrosis as determined by non-invasive tests in patients with cardiometabolic risk factors without other causes of steatosis, notably, alcohol consumption of  $> 20$  g/day for women and  $> 30$  g/day for men. Specifically, fibrosis can be estimated using imaging and/or blood-based non-invasive tests with reasonable to high accuracy.<sup>22</sup> There are no FDA-approved non-invasive tests to diagnose MASH with stage F2 to F3 fibrosis or to monitor the response to pharmacotherapy.<sup>21</sup> Although liver biopsy is not typically recommended for fibrosis staging in current clinical practice, histologic exam remains the gold standard to quantify fibrosis if performed previously (historical biopsy obtained reasonably recently, e.g., within 3 years).<sup>21</sup> Since non-invasive tests are more readily available than liver biopsy, it is recommended that more current data (e.g., within 6 to 12 months) be utilized to determine patients who are appropriate candidates for treatment. Three non-invasive tests are supported in the 2025 guidance: vibration-controlled transient elastography (VCTE), magnetic resonance elastography (MRE), and the blood-based biomarker Enhanced Liver Fibrosis™ (ELF) score.<sup>22</sup> For the blood-based ELF score, a cutoff of 9.2 provides optimal sensitivity and specificity for detecting  $\geq$  F2 fibrosis. To guide the selection of appropriate candidates for treatment with MASH therapy, the ELF score range of 9.2 to 10.5 is recommended. In the ESSENCE trial, this range identified approximately 50% of patients with F2 or F3 fibrosis. While the range serves as guidance, in those with values outside of the upper range, exclusion of cirrhosis and portal hypertension should be considered using another non-invasive test such as VCTE or MRE. Wegovy is not indicated in patients with cirrhotic (F4) MASH.

### *Sleep Apnea*

The American Academy of Sleep Medicine (2017) recommends that diagnostic testing for OSA be performed in combination with a comprehensive sleep evaluation.<sup>10</sup> Polysomnography is the gold standard test for the diagnosis of OSA in adults in whom there is concern for OSA based on the sleep evaluation. In some cases, and within the appropriate context, the use of home sleep apnea test as the initial sleep study may be acceptable, however, polysomnography should be used when home sleep apnea test results do not provide satisfactory posttest probability of confirming or ruling out OSA.

Available treatment guidelines for OSA do not specifically mention the GLP-1 agonists. The American Thoracic Society clinical practice guideline on the role of weight management in the treatment with adults with OSA (2018) recommends that patients with OSA who are overweight or obese (BMI  $\geq 25$  kg/m<sup>2</sup>) participate in comprehensive lifestyle intervention that includes a

reduced calorie diet, exercise/increased physical activity, and behavioral counseling.<sup>11</sup> For patients with OSA and a BMI  $\geq 27$  kg/m<sup>2</sup> who have not had an improvement in weight despite a comprehensive weight-loss lifestyle program, and have no contraindications (no active CV disease), evaluation for anti-obesity medication is suggested. The weight-loss goal in patients with overweight or obesity with OSA should be  $\geq 7\%$  to  $10\%$  of total body weight. The AACE Consensus Statement: Algorithm for the Evaluation and Treatment of Adults with Obesity/ABCD (2025 update) recommends a weight loss target of  $7\%$  to  $10\%$  in patients with OSA and ABCD/obesity;  $> 10\%$  weight loss is noted to result in additional benefit.<sup>23</sup>

Clinical success in OSA has been described by several publications. The American Academy of Sleep Medicine (2019) cites a clinically significant threshold of  $\geq 15$  events/hour (on AHI)<sup>12</sup> and a clinical practice guideline for the treatment of OSA and snoring with oral appliance therapy (2015) from the American Academy of Sleep Medicine and American Academy of Dental Sleep Medicine<sup>13</sup> notes that treatment success is usually defined as a reduction in the AHI to a specific level (e.g., post-treatment AHI  $< 5$  events/hour, or a  $> 50\%$  reduction in AHI). Of note, a meta-analysis on the impact of weight reduction on AHI reported that weight reduction in patients with obesity and OSA was associated with improvements in the severity of OSA. A BMI reduction of  $20\%$  was associated with an AHI reduction of  $57\%$ ; further weight reduction beyond  $20\%$  in BMI was associated with a smaller effect on AHI.<sup>14</sup>

## Coverage Policy

### Policy Statement

Prior Authorization is required for benefit coverage of Foundayo, liraglutide (Saxenda, generic), Wegovy injection, Wegovy tablet and Zepbound. Of note, this policy targets Foundayo, liraglutide (Saxenda, generic), Wegovy, and Zepbound; other glucagon-like peptide-1 agonists which do not carry an FDA-approved indication for weight loss are not targeted in this policy. In the clinical criteria, as appropriate, an asterisk (\*) is noted next to the specified gender. In this context, the specified gender is defined as follows: males are defined as individuals with the biological traits of a male, regardless of the individual's gender identity or gender expression; females are defined as individuals with the biological traits of a female, regardless of the individual's gender identity or gender expression. Because of the specialized skills required for evaluation and diagnosis of patients treated with Wegovy injection for metabolic dysfunction-associated steatohepatitis (MASH)/non-alcoholic steatohepatitis (NASH) as well as the monitoring required for adverse events and long-term efficacy, approval requires Wegovy injection for MASH/NASH to be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days.

**Documentation:** Documentation is required for use of Saxenda for weight loss and Wegovy injection for MASH/NASH as noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information.

**Glucagon-like peptide-1 (GLP-1) receptor agonists are considered medically necessary when ONE of the following is met (I, II, III, IV, or V):**

**I. Liraglutide (Saxenda, generic) is considered medically necessary when ONE of the following is met (1 or 2):**

#### **FDA-Approved Indications**

- 1. Weight Loss in an Adult with Obesity or is Overweight.** Approve for the duration noted if the patient meets ONE of the following (A or B):
- A) Initial Therapy.** Approve for 4 months if the patient meets ALL of the following (i, ii, iii, iv and v):
- i.** Patient is  $\geq 18$  years of age; AND
  - ii.** Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND
  - iii.** Patient meets ONE of the following (a or b):
    - a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
    - b)** Patient meets BOTH of the following [(1) and (2)]:
      - (1)**At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND
      - (2)**At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
  - iv.** The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; AND
  - v.** Preferred product criteria is met for the product(s) as listed in the below table(s); OR
- B) Patient is Currently Receiving liraglutide (Saxenda, generic).** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):  
Note: For a patient who has not completed 4 months of initial therapy, refer to Initial Therapy criteria above.
- i.** Patient is  $\geq 18$  years of age; AND
  - ii.** Patient meets ONE of the following (a or b):
    - a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
    - b)** Patient meets BOTH of the following [(1) and (2)]:
      - (1)**At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND
      - (2)**At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-

1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has lost  $\geq 4\%$  of baseline body weight; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**2. Weight Loss in a Pediatric Patient with Obesity.** Approve for the duration noted if the patient meets ONE of the following (A or B):

**A) Initial Therapy.** Approve for 4 months if the patient meets ALL of the following (i, ii, iii, iv and v):

i. Patient is  $\geq 12$  years of age and  $< 18$  years of age; AND

ii. Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND

iii. At baseline, patient had a BMI  $\geq 95^{\text{th}}$  percentile for age and sex; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; AND

v. Preferred product criteria is met for the product(s) as listed in the below table(s);  
OR

**B) Patient is Currently Receiving liraglutide (Saxenda, generic).** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 4 months of initial therapy, refer to Initial Therapy criteria above.

i. Patient is  $\geq 12$  years of age and  $< 18$  years of age; AND

ii. At baseline, patient had a BMI  $\geq 95^{\text{th}}$  percentile for age and sex; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has had a reduction in BMI of  $\geq 1\%$  from baseline; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**Employer Plans:**

Product	Criteria
<b>Saxenda</b> (liraglutide subcutaneous injection)	The patient has tried the bioequivalent generic product, <b><u>liraglutide subcutaneous injection (generic for Saxenda)</u></b> , AND cannot take due to a formulation difference in the inactive ingredient(s) [e.g., difference in dyes, fillers, preservatives] between the brand and the bioequivalent generic product which, per the prescriber, would result in a significant allergy or serious adverse reaction. <b>[documentation required]</b>

**II. Foundayo is considered medically necessity when the following is met:**

**FDA-Approved Indication**

**1. Weight Loss in an Adult with Obesity or is Overweight.** Approve for the duration noted if the patient meets ONE of the following (A or B):

**A) Initial Therapy.** Approve for 8 months if the patient meets ALL of the following (i, ii, iii, and iv):

**i.** Patient is  $\geq 18$  years of age; AND

**ii.** Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND

**iii.** Patient meets ONE of the following (a or b):

**a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**b)** Patient meets BOTH of the following [(1) and (2)]:

**(1)**At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND

**(2)**At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**iv.** The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

**B) Patient is Currently Receiving Foundayo.** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 8 months of initial therapy, refer to Initial Therapy criteria above.

**i.** Patient is  $\geq 18$  years of age; AND

**ii.** Patient meets ONE of the following (a or b):

**a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**b)** Patient meets BOTH of the following [(1) and (2)]:

**(1)**At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND

**(2)**At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has lost  $\geq 5\%$  of baseline body weight; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**III. Wegovy injection is considered medically necessary when ONE of the following is met (1, 2, 3 or 4):**

**FDA-Approved Indications**

**1. Weight Loss in an Adult with Obesity or is Overweight.** Approve for the duration noted if the patient meets ONE of the following (A or B):

**A) Initial Therapy.** Approve for 8 months if the patient meets ALL of the following (i, ii, iii, and iv):

i. Patient is  $\geq 18$  years of age; AND

ii. Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND

iii. Patient meets ONE of the following (a or b):

**a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**b)** Patient meets BOTH of the following [(1) and (2)]:

**(1)** At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND

**(2)** At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

**B) Patient is Currently Receiving Wegovy injection or Wegovy tablet.** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 8 months of initial therapy with Wegovy injection and/or Wegovy tablet, refer to Initial Therapy criteria above.

i. Patient is  $\geq 18$  years of age; AND

ii. Patient meets ONE of the following (a or b):

**a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**b)** Patient meets BOTH of the following [(1) and (2)]:

**(1)**At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND

**(2)**At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**iii.** Patient has lost  $\geq 5\%$  of baseline body weight; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**iv.** The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**2. Weight Loss in a Pediatric Patient with Obesity.** Approve for the duration noted if the patient meets ONE of the following (A or B):

**A) Initial Therapy.** Approve for 8 months if the patient meets ALL of the following (i, ii, iii, and iv):

**i.** Patient is  $\geq 12$  years of age and  $< 18$  years of age; AND

**ii.** Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND

**iii.** At baseline, patient had a BMI  $\geq 95^{\text{th}}$  percentile for age and sex; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**iv.** The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

**B) Patient is Currently Receiving Wegovy injection or Wegovy tablet.** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 8 months of initial therapy with Wegovy injection and/or Wegovy tablet, refer to Initial Therapy criteria above.

**i.** Patient is  $\geq 12$  years of age and  $< 18$  years of age; AND

**ii.** At baseline, patient had a BMI  $\geq 95^{\text{th}}$  percentile for age and sex; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**iii.** Patient has had a reduction in BMI of  $\geq 1\%$  from baseline; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

- iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**3. Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease with Overweight or Obesity.** Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):

- A)** Patient is  $\geq 18$  years of age; AND
- B)** At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
- C)** Patient meets ONE of the following (i, ii, or iii):
  - i.** Patient has had a prior myocardial infarction; OR
  - ii.** Patient has had a prior stroke; OR  
Note: This does not include a transient ischemic attack (TIA).
  - iii.** Patient has a history of symptomatic peripheral arterial disease as evidenced by ONE of the following [a, b, or c]:
    - a)** Intermittent claudication with ankle-brachial index  $< 0.85$ ; OR
    - b)** Peripheral arterial revascularization procedure; OR
    - c)** Amputation due to atherosclerotic disease; AND
- D)** According to the prescriber, the medication will be used in combination with optimized pharmacotherapy for established cardiovascular disease; AND
- E)** The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**4. Metabolic Dysfunction-Associated Steatohepatitis (MASH)/Non-Alcoholic Steatohepatitis (NASH).** Approve for 1 year if the patient meets the ONE of the following (A or B):

- A)** Initial Therapy: Approve if the patient meets ALL of the following (i, ii, iii, iv, v, vi, vi, and vii):
  - i.** Patient is  $\geq 18$  years of age; AND
  - ii.** Patient does not have cirrhosis (F4); AND
  - iii.** Patient has stage F2 or F3 fibrosis prior to initiating treatment with Rezdifra, Wegovy injection, or Wegovy tablet **[documentation required]** identified by ONE of the following (a, b, c, or d):
    - a)** Liver biopsy performed within the 3 years preceding treatment with Rezdifra, Wegovy injection, or Wegovy tablet **[documentation required]**; OR
    - b)** Vibration-controlled transient elastography (VCTE) performed within the 6 months preceding treatment with Rezdifra, Wegovy injection, or Wegovy tablet **[documentation required]**; OR
    - c)** Magnetic resonance elastography (MRE) performed within the 6 months preceding treatment with Rezdifra, Wegovy injection, or Wegovy tablet **[documentation required]**; OR
    - d)** Enhanced Liver Fibrosis™ (ELF) test performed within the 6 months preceding treatment with Rezdifra, Wegovy injection, or Wegovy tablet **[documentation required]** with a score of  $\geq 9.2$  to  $\leq 10.5$  **[documentation required]**; AND
  - iv.** According to the prescriber, the patient has ONE or more of the following metabolic risk factors that are managed according to standard of care (a, b, c, d, e):
    - a)** Central obesity;

- b)** Hypertriglyceridemia;
  - c)** Reduced high-density lipoprotein cholesterol;
  - d)** Hypertension;
  - e)** Elevated fasting plasma glucose indicative of diabetes or pre-diabetes; AND
  - v.** According to the prescriber, patient meets ONE of the following (a or b):
    - a)** Female\* patient: Alcohol consumption is < 20 grams/day; OR  
Note: One standard drink (or one alcoholic drink equivalent) contains roughly 14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.
    - b)** Male\* patient: Alcohol consumption < 30 grams/day; AND  
Note: One standard drink (or one alcoholic drink equivalent) contains roughly 14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.
  - vi.** The medication will be used in combination with appropriate diet and exercise therapy; AND
  - vii.** The medication is prescribed by or in consultation with an endocrinologist, gastroenterologist, or hepatologist; OR
- B) Patient is Currently Receiving Wegovy injection or Wegovy tablet: Approve if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):**  
Note: For a patient who has received < 1 year of therapy with Wegovy injection and/or Wegovy tablet or who is restarting therapy, refer to Initial Therapy criteria above.
- i.** Patient has completed ≥ 1 year of therapy with Wegovy injection and/or Wegovy tablet AND according to the prescriber, the patient has not had worsening of fibrosis or MASH/NASH; AND  
Note: This applies to a patient starting their second (or more) year of therapy with Wegovy injection and/or Wegovy tablet (i.e., the patient has already completed 1 year or more of therapy with Wegovy injection and/or tablet).
  - ii.** According to the prescriber, patient does not have cirrhosis (F4); AND
  - iii.** According to the prescriber, metabolic risk factors are managed according to standard of care; AND
  - iv.** According to the prescriber, patient meets ONE of the following (a or b):
    - a)** Female\* patient: Alcohol consumption is < 20 grams/day; OR  
Note: One standard drink (or one alcoholic drink equivalent) contains roughly 14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.
    - b)** Male\* patient: Alcohol consumption < 30 grams/day; AND  
Note: One standard drink (or one alcoholic drink equivalent) contains roughly 14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.
  - v.** The medication will be used in combination with appropriate diet and exercise therapy; AND
  - vi.** The medication is prescribed by or in consultation with an endocrinologist, gastroenterologist, or hepatologist.

\*Refer to the Policy Statement

**IV. Wegovy tablet is considered medically necessary when ONE of the following is met (1 or 2):**

## FDA-Approved Indications

**1. Weight Loss in an Adult with Obesity or is Overweight.** Approve for the duration noted if the patient meets ONE of the following (A or B):

**A) Initial Therapy.** Approve for 8 months if the patient meets ALL of the following (i, ii, iii, and iv):

- i.** Patient is  $\geq 18$  years of age; AND
- ii.** Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND
- iii.** Patient meets ONE of the following (a or b):
  - a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
  - b)** Patient meets BOTH of the following [(1) and (2)]:
    - (1)**At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND
    - (2)**At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
- iv.** The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

**B) Patient is Currently Receiving Wegovy tablet or Wegovy injection.** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 8 months of initial therapy with Wegovy tablet and/or Wegovy injection, refer to Initial Therapy criteria above.

- i.** Patient is  $\geq 18$  years of age; AND
- ii.** Patient meets ONE of the following (a or b):
  - a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
  - b)** Patient meets BOTH of the following [(1) and (2)]:
    - (1)**At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND
    - (2)**At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-

1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has lost  $\geq 5\%$  of baseline body weight; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**2. Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease with Overweight or Obesity.** Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):

**A)** Patient is  $\geq 18$  years of age; AND

**B)** At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**C)** Patient meets ONE of the following (i, ii, or iii):

**i.** Patient has had a prior myocardial infarction; OR

**ii.** Patient has had a prior stroke; OR

Note: This does not include a transient ischemic attack (TIA).

**iii.** Patient has a history of symptomatic peripheral arterial disease as evidenced by ONE of the following [a, b, or c]:

**a)** Intermittent claudication with ankle-brachial index  $< 0.85$ ; OR

**b)** Peripheral arterial revascularization procedure; OR

**c)** Amputation due to atherosclerotic disease; AND

**D)** According to the prescriber, the medication will be used in combination with optimized pharmacotherapy for established cardiovascular disease; AND

**E)** The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**V. Zepbound is considered medically necessary when ONE of the following is met (1 or 2):**

#### **FDA-Approved Indications**

**1. Weight Loss in an Adult with Obesity or is Overweight.** Approve for the duration noted if the patient meets ONE of the following (A or B):

**A) Initial Therapy.** Approve for 8 months if the patient meets ALL of the following (i, ii, iii, and iv):

**i.** Patient is  $\geq 18$  years of age; AND

**ii.** Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND

**iii.** Patient meets ONE of the following (a or b):

**a)** At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

**c)** Patient meets BOTH of the following [(1) and (2)]:

**(1)** At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND

(2) At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

**B) Patient is Currently Receiving Zepbound.** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 8 months of initial therapy, refer to Initial Therapy criteria above.

i. Patient is  $\geq 18$  years of age; AND

ii. Patient meets ONE of the following (a or b):

a) At baseline, patient had a BMI  $\geq 35$  kg/m<sup>2</sup>; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

b) Patient meets BOTH of the following [(1) and (2)]:

(1) At baseline, patient had a BMI  $\geq 27$  kg/m<sup>2</sup>; AND

(2) At baseline, patient had, or patient currently has, at least TWO of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has lost  $\geq 5\%$  of baseline body weight; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

**2. Obstructive Sleep Apnea, Moderate to Severe, in a Patient with Obesity.** Approve for 1 year if the patient meets ONE of the following (A or B):

**A) Initial Therapy.** Approve if the patient meets ALL of the following (i, ii, iii, iv, and v):

i. Patient is  $\geq 18$  years of age; AND

ii. At baseline, patient had a BMI  $\geq 30$  kg/m<sup>2</sup>; AND

- Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
- iii. Patient has had a sleep study that shows BOTH of the following (a and b):
    - a) Patient has been diagnosed with moderate to severe obstructive sleep apnea; AND
    - b) Patient has an apnea-hypopnea index  $\geq$  15 events per hour; AND  
Note: A diagnosis of moderate obstructive sleep apnea is an apnea-hypopnea index of  $\geq$  15 events per hour and a diagnosis of severe sleep apnea is an apnea-hypopnea index  $\geq$  30 events per hour. The apnea-hypopnea index is the number of apneas and hypopneas during 1 hour of sleep.
  - iv. The patient does NOT meet either of the following (a or b):  
Note: A patient who has one or more of the following conditions/diagnoses below is not approved.
    - a) Central sleep apnea; OR
    - b) Cheyne Stokes respiration; AND
  - v. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

**B) Patient is Currently Receiving Zepbound.** Approve if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 1 year of initial therapy, refer to Initial Therapy criteria above.

- i. Patient is  $\geq$  18 years of age; AND
- ii. At baseline, patient had a BMI  $\geq$  30 kg/m<sup>2</sup>; AND  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
- iii. Patient has completed  $\geq$  1 year of therapy with Zepbound AND the patient meets BOTH of the following (a and b):
  - a) Patient has lost  $\geq$  10% of baseline body weight; AND  
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Foundayo, liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
  - b) According to the prescriber, patient has stability in obstructive sleep apnea signs or symptoms; AND  
Note: Examples of signs or symptoms of obstructive sleep apnea include but are not limited to snoring, excessive daytime sleepiness, fatigue.
- iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

### Conditions Not Covered

**Foundayo, Liraglutide (Saxenda, generic), Wegovy injection, Wegovy tablet, and Zepbound for any other use are considered not medically necessary, including the following (this list may not be all inclusive; criteria will be updated as new published data are available):**

- 1. Concomitant Use with other Glucagon-Like Peptide-1 (GLP-1) Agonists or GLP-1/ Glucose-Dependent Insulinotropic Polypeptide (GIP) Receptor Agonists.** The GLP-

1 agonists and the GLP-1/GIP agonist should not be combined with each other or with any other GLP-1 agonists or GLP-1/GIP agonist.<sup>1,2,4,25</sup> There are other GLP-1 and GLP-1/GIP products not included in this policy that are FDA-approved for type 2 diabetes, and not for chronic weight management.

**Note:** Examples of other GLP-1 agonists include but are not limited to exenatide SC injection, Ozempic (semaglutide tablets and SC injection), Rybelsus (semaglutide tablets), Trulicity (dulaglutide SC injection), and liraglutide SC injection (Victoza, generic). An example of a GLP-1/GIP agonist is Mounjaro (tirzepatide SC injection).

## References

1. Wegovy<sup>®</sup> tablet and subcutaneous injection [prescribing information]. Plainsboro, NJ: Novo Nordisk; March 2026.
2. Saxenda<sup>®</sup> subcutaneous injection [prescribing information]. Plainsboro, NJ: Novo Nordisk; May 2025.
3. Hampl SE, Hassink SG, Skinner AC, et al. Clinical Practice Guideline for the Evaluation and Treatment of Children and Adolescents with Obesity. *Pediatrics*. 2023;151(2):e2022060640.
4. Zepbound<sup>®</sup> subcutaneous injection [prescribing information]. Indianapolis, IN: Eli Lilly; April 2025.
5. Lincoff AM, Brown-Frandson K, Colhoun HM, et al; for the SELECT Trial Investigators. Semaglutide and cardiovascular outcomes in obesity without diabetes. *N Engl J Med*. 2023;389(24):2221-2232
6. Lingvay I, Brown-Frandson K, Colhoun HM et al. Semaglutide for cardiovascular event reduction in people with overweight or obesity: SELECT study baseline characteristics. *Obesity*. 2023;31(1):111-122.
7. Wilding JPH, Batterham RL, Calanna S, et al; STEP 1 Study Group. Once-weekly semaglutide in adults with overweight or obesity. *N Engl J Med*. 2021;384(11):989.
8. Weghuber D, Barrett T, Barrientos-Pérez M, et al; STEP TEENS Investigators. Once-weekly semaglutide in adolescents with obesity. *N Engl J Med*. 2022;387(24):2245-2257.
9. Malhorta A, Grunstein RR, Fietze I, et al; for the SURMOUNT-OSA Investigators. Tirzepatide for the treatment of obstructive sleep apnea and obesity. *N Engl J Med*. 2024;391(13):1193-1205.
10. Kapur VK, Auckley DH, Chowdhuri S, et al. Clinical practice guideline for diagnostic testing for adult obstructive sleep apnea: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2017;13(3):479-504.
11. Hugel DW, Patel SR, Ahasic AM, et al; on behalf of the American Thoracic Society Assembly on Sleep and Respiratory Neurology. The role of weight management in the treatment of adult obstructive sleep apnea. *Am J Respir Crit Care Med*. 2018;198(6): e70-e87.
12. Ramar K, Dort LC, Katz SG et al. Clinical practice guideline for the treatment of obstructive sleep apnea and snoring with oral appliance therapy: an update for 2015. An American Academy of Sleep Medicine and American Academy of Dental Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med*. 2015;11(7):773-827.
13. Patil SP, Ayappa IA, Caples SM, et al. Treatment of adult obstructive sleep apnea with positive airway pressure: An American Academy of Sleep Medicine Systematic Review, Meta-analysis, and GRADE Assessment. *J Clin Sleep Med*. 2019;15(2):301-334.
14. Malhorta A, Heilman CR, Banerjee, et al. Weight reduction and the impact on apnea-hypopnea index: a systematic meta-analysis. *Sleep Medicine*. 2023;121:26-31.
15. Jasterboff AM, Aronne LJ, Ahmad NN, et al; for the SURMOUNT-1 Investigators. Tirzepatide once weekly for the treatment of obesity. *N Engl J Med*. 2022;387(3):205-216.
16. Garvey TW, Frias JP, Jasterboff, et al; for the SURMOUNT-2 Investigators. Tirzepatide once weekly for the treatment of obesity in people with type 2 diabetes (SURMOUNT-2): a double-

- blind, randomized, multicenter, placebo-controlled, phase 3 trial. *Lancet*. 2023;402(10402):613-6262.
17. Wadden TA, Chao AM, Machineni, et al. Tirzepatide after intensive lifestyle intervention in adults with overweight or obesity: The SURMOUNT-3 phase 3 trial. *Nature Med*. 2023;29(11):2909-2918.
  18. Wadden TA, Hollander P, Klein S, et al; on behalf of the NN8022-1923 Investigators. Weight maintenance and additional weight loss with liraglutide after low-calorie-diet-induced weight loss: The SCALE Maintenance randomized study. *Int J Obes*. 2013;37:1443-1451.
  19. Newsome PN, Sanyal AJ, Engerbretsen KA, et al. Semaglutide 2.4 mg in participants with metabolic dysfunction-associated steatohepatitis: baseline characteristics and design of the phase 3 ESSENCE trial. *Aliment Pharmacol Ther*. 2024;60(11-12):1525-1533.
  20. Sanyal AJ, Newsome PN, Kliers I, et al. Phase 3 trial of semaglutide in metabolic dysfunction-associated steatohepatitis. *N Engl J Med*. 2025;392(21):2089-2099.
  21. Chen VL, Morgan TR, Rotman Y, et al. Resmetirom therapy for metabolic dysfunction-associated steatotic liver disease: October 2024 updates to AASLD Practice Guidance. *Hepatology*. 2025;81(1):312-320.
  22. Bansal MB, Patton H, Morgan TR, et al. Semaglutide therapy for metabolic dysfunction-associated steatohepatitis: November 2025 updates to AASLD practice guidance. *Hepatology*. 2025 Nov 7. [Online ahead of print].
  23. Nadolsky K, Garvey WT, Agarwal M, et al. American Association of Clinical Endocrinology consensus statement: algorithm for the evaluation and treatment of adults with obesity/adiposity-based chronic disease — 2025 Update. *Endocrine Practice*. 2025;31:1351–1394.
  24. American Diabetes Association Professional Practice Committee for Obesity. Pharmacologic treatment of obesity in adults: Standards of care in overweight and obesity. *DOCM Care*. 13 Jan 2026 [Online ahead of print].
  25. Foundayo™ tablets [prescribing information]. Indianapolis, IN: Eli Lilly; April 2026.

## Revision Details

Summary of Changes	Review Date	Effective Date
New policy	04/24/2025	07/01/2025
<p><u>Wegovy:</u>  <b>Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease who is Either Obese or Overweight.</b>  <u>Initial Therapy.</u> For the requirement that a patient has had a prior stroke, a note was added that a to clarify that this does not include a transient ischemic attack (TIA).</p> <p><u>Zepbound:</u>  <b>Obstructive Sleep Apnea, Moderate to Severe, in a Patient with Obesity.</b>  <u>Initial Therapy.</u> The requirement that a patient had a sleep study was modified to remove the timeframe that the sleep study was within the past 1 year. A patient is still required to have a sleep study.</p> <p>Conditions Not Covered:</p>	06/12/2025	08/01/2025

<p><b>Concomitant Use with Other Medications FDA-Approved for Weight Loss.</b> This condition not recommended for approval was clarified to state that concomitant use with other medications FDA-approved for weight loss is not recommended. Previously, the requirement did not specify medications were "FDA-approved" for weight loss. The note with examples of weight loss medications was updated to reflect product availability for medications FDA-approved for weight loss.</p> <p><b>Concomitant Use with Glucagon-Like Peptide-1 (GLP-1) Agonists or GLP-1/ Glucose-Dependent Insulinotropic Polypeptide (GIP) Agonists.</b> The note was updated to reflect availability for other GLP-1 or GLP-1/GIP agonists.</p>		
<p><u>Saxenda:</u>  <b>Weight Loss in an Adult with Overweight or Obesity.</b> This condition of approval was modified to add "with obesity or is overweight".  <b>Weight Loss in a Pediatric Patient with Obesity.</b> This condition of approval was modified to add "with obesity".</p> <p><u>Wegovy:</u>  <b>Weight Loss in an Adult with Overweight or Obesity.</b> This condition of approval was modified to add "with obesity or is overweight".  <b>Weight Loss in a Pediatric Patient with Obesity.</b> This condition of approval was modified to add "with obesity".  <b>Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease in a Patient with Obesity or Overweight.</b> This condition of approval was re-worded from "in an overweight or obese patient" to "in a patient with obesity or is overweight".</p> <p><u>Zepbound:</u>  <b>Weight Loss in an Adult with Overweight or Obesity.</b> This condition of approval was modified to add "with obesity or is overweight".</p>	07/17/2025	09/15/2025
<p>Liraglutide, generic to Saxenda was added to the policy.</p> <p><b>Policy Statement:</b> The following was added: In clinical criteria, as appropriate, an asterisk (*) is noted next to the specified gender. In this context, the specified gender is defined as follows: males are defined as individuals with the biological traits of a male, regardless of the individual's gender identity or gender expression; females are defined as individuals with the biological traits of a female, regardless of the individual's gender identity or gender expression. Because of the specialized skills required for evaluation and diagnosis of patients treated with Wegovy as well as the</p>	11/06/2025	12/15/2025

monitoring required for adverse events and long-term efficacy, approval requires Wegovy to be prescribed by or in consultation with a physician who specializes in the condition being treated. The Policy Statement was updated as follows to address the availability of liraglutide, generic to Saxenda: Prior Authorization is recommended for prescription benefit coverage of liraglutide (Saxenda, generic), Wegovy, and Zepbound. Of note, this policy targets liraglutide (Saxenda, generic), Wegovy, and Zepbound; other glucagon-like peptide-1 agonists that do not carry an FDA-approved indication for weight loss are not targeted in this policy.

**Documentation:** A requirement for documentation was added for the use of Wegovy for MASH. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information.

Wegovy:

**Metabolic Dysfunction-Associated Steatohepatitis (MASH)/Non-Alcoholic Steatohepatitis (NASH).** A new condition of coverage was added to FDA-Approved Indications.

Conditions Not Covered:

**Concomitant Use with Other Medications FDA-Approved for Weight Loss.** This condition not recommended for approval was removed.

Liraglutide (Saxenda, generic), Wegovy, and Zepbound:  
**Weight Loss in an Adult with Overweight or Obesity:**  
Initial Therapy and Patient is Continuing on Therapy: The notes that define baseline were updated to include liraglutide, generic to Saxenda.

Liraglutide (Saxenda, generic) and Wegovy:  
**Weight Loss in a Pediatric Patient with Obesity:**  
Initial Therapy and Patient is Continuing on Therapy: The notes that define baseline were updated to include liraglutide, generic to Saxenda.

Saxenda

**Weight Loss in an Adult with Obesity or is Overweight.**  
Initial therapy

**Added** preferred product requirements.

**Weight Loss in a Pediatric Patient with Obesity.**  
Initial therapy

**Added** preferred product requirements.

<p><b>Updated</b> the policy statement.  <b>Changed</b> each "Patient is Continuing Therapy" to "Patient is Currently Receiving"</p> <p><u>Wegovy</u>  <b>Metabolic Dysfunction-Associated Steatohepatitis (MASH)/Non-Alcoholic Steatohepatitis (NASH).</b>  <u>Initial Therapy.</u> The requirement that the patient does not have cirrhosis was clarified that the patient does not have cirrhosis "(F4)". Criteria for the diagnosis of MASH/NASH were updated such that the patient must have documentation of stage F2 or F3 fibrosis prior to initiating treatment with Rezdifra or Wegovy confirmed by ONE of the following: Liver biopsy performed within 3 years preceding treatment with Rezdifra or Wegovy <b>[documentation required]</b>, vibration-controlled elastography (VCTE) performed within 6 months preceding treatment with Rezdifra or Wegovy <b>[documentation required]</b>, magnetic resonance imaging (MRE) performed within 6 months preceding treatment with Rezdifra or Wegovy <b>[documentation required]</b>, or Enhanced Liver Fibrosis (ELF) test performed within 6 months preceding treatment with Rezdifra or Wegovy <b>[documentation required]</b> with a score of <math>\geq 9.2</math> to <math>\leq 10.5</math> <b>[documentation required]</b>. Previously, the diagnosis of MASH/NASH was confirmed by either a liver biopsy within 3 years preceding treatment with Wegovy <b>[documentation required]</b> that showed a non-alcoholic fatty liver disease activity score of <math>\geq 4</math> with a score of <math>\geq 1</math> in steatosis, ballooning, and lobular inflammation <b>[documentation required]</b> OR an imaging exam (i.e., elastography, computed tomography, or magnetic resonance imaging) performed within 6 months preceding treatment with Wegovy <b>[documentation required]</b>. The separate criterion that the patient have stage F2 or F3 fibrosis <b>[documentation required]</b> was removed (this is part of the updated criterion outlined above; patients must still have documentation of F2 or F3 fibrosis). Reference to prior to initiating therapy throughout criteria were updated to include Rezdifra (i.e., prior to initiating treatment with Rezdifra or Wegovy); previously only Wegovy.  <u>Patient is Currently Receiving Wegovy.</u> The criterion requiring that according to the prescriber the patient has not progressed to stage F4 (cirrhosis) was modified to state, according to the prescriber, patient does not have cirrhosis (F4).  Zepbound  <b>Obstructive Sleep Apnea, Moderate to Severe, in a Patient with Obesity.</b></p>	<p>12/18/2025</p>	<p>02/15/2026</p>
--	-------------------	-------------------

<p><u>Initial Therapy</u>. The criterion excluding coverage of a patient with central sleep apneas was modified to remove the additional requirement that the percent of central apneas/hypopneas is <math>\geq 50\%</math>.</p>		
<p>Wegovy tablet was added to the policy; new criteria were created.</p> <p>Wegovy injection  <b>Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease who is Either Obese or Overweight.</b>  <u>Patient is Currently Receiving Wegovy injection</u>. The note that baseline body mass index refers to baseline prior to Wegovy injection was updated to also include Wegovy tablet.</p>	02/12/2026	03/01/2026
<p>Foundayo was added to the Policy; new criteria were created.</p> <p><b>Policy Statement:</b> Foundayo was added the list of products to which the Policy applies. The statement was also updated to reflect the availability of Wegovy tablet and Wegovy injection.</p> <p>Throughout the policy, Foundayo was added as an example of a glucagon-like peptide-1 (GLP-1) receptor agonist.</p> <p><u>Wegovy injection</u>  <b>Weight Loss in an Adult with Obesity or is Overweight.</b>  The initial approval was changed to 8 months. The corresponding Note for a patient currently receiving Wegovy tablet or Wegovy injection was updated to reflect the change to initial approval. Wegovy tablet was added to the following: <u>Patient is Currently Receiving Wegovy injection or Wegovy tablet</u>. The following note was updated to include Wegovy injection and Wegovy tablet: For a patient who has not completed 7 months of initial therapy with Wegovy injection and/or Wegovy tablet, refer to Initial Therapy criteria above.  <b>Weight Loss in a Pediatric Patient with Obesity.</b> The initial approval was changed to 8 months. The corresponding Note for a patient currently receiving Wegovy tablet or Wegovy injection was updated to reflect the change to initial approval. Wegovy tablet was added to the following: <u>Patient is Currently Receiving Wegovy injection or Wegovy tablet</u>. The following note was updated to include Wegovy injection and Wegovy tablet: For a patient who has not completed 7 months of initial therapy with Wegovy injection and/or Wegovy tablet, refer to Initial Therapy criteria above.  <b>Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease with Overweight or Obesity.</b> <u>Initial Therapy</u> and <u>Patient is Currently Receiving Wegovy injection</u> criteria were combined</p>	04/16/2026	04/30/2026

(previously each a 1 year approval); the approval duration is 1 year. The requirement that the patient has a current body mass index (BMI) of  $\geq 27 \text{ kg/m}^2$  (previously for Initial Therapy, only) was revised that at baseline, the patient had a BMI  $\geq 27 \text{ kg/m}^2$ . The note for baseline BMI was revised to state that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound) [previously Wegovy injection or Wegovy tablet, Patient is Currently Receiving Wegovy injection or Wegovy tablet].

**Metabolic Dysfunction-Associated Steatohepatitis (MASH)/Non-Alcoholic Steatohepatitis (NASH).** Initial

Therapy. Reference to Wegovy throughout criteria was updated to specify Wegovy injection. Wegovy tablet for the requirements related to identification of stage F2 or F3 fibrosis prior to initiating treatment with Rezdiffra, Wegovy injection, or Wegovy tablet [**documentation required**].

Wegovy tablet was added to the following: Patient is Currently Receiving Wegovy injection or Wegovy tablet. The following note was updated to include Wegovy injection and Wegovy tablet: For a patient who has received  $< 1$  year of therapy with Wegovy injection and/or Wegovy tablet, or who is restating therapy, refer to Initial Therapy. For the following requirement (and associated Note), reference to Wegovy was updated to specify Wegovy injection and to add Wegovy tablet: Patient has completed  $\geq 1$  year of therapy with Wegovy injection and/or Wegovy tablet AND according to the prescriber, the patient has not had worsening of fibrosis or MASH/NASH.

Wegovy tablet

**Weight Loss in an Adult with Obesity or is Overweight.**

The initial approval was changed to 8 months. The corresponding Note for a patient currently receiving Wegovy tablet or Wegovy injection was updated to reflect the change to initial approval.

Initial Therapy. The approval duration was changed to 7 months (previously 6 months). Wegovy injection was added to Patient is Currently Receiving Wegovy tablet or Wegovy injection. The following note was updated to include Wegovy injection and Wegovy tablet, and to reflect the 7-month initial approval duration: For a patient who has not completed 7 months of initial therapy with Wegovy injection and/or Wegovy tablet, refer to Initial Therapy criteria above.

**Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease with Overweight or Obesity.** The same changes were made as Wegovy injection; refer to Wegovy injection *Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease with Overweight or Obesity*, above.

<p><u>Zepbound</u>  <b>Obstructive Sleep Apnea, Moderate to Severe, in a Patient with Obesity.</b> <u>Initial Therapy.</u> The requirement that the patient has a current BMI <math>\geq 30</math> kg/m<sup>2</sup> was revised that at baseline, patient had a BMI <math>\geq 30</math> kg/m<sup>2</sup>. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound). <u>Patient is Currently Receiving Zepbound.</u> The notes that define baseline were updated that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound); previously, prior to Zepbound.</p>		
--	--	--

The policy effective date is in force until updated or retired.

---

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2026 The Cigna Group.