



## Drug Coverage Policy

Effective Date .....2/15/2026  
Coverage Policy Number.....IP0768  
Policy Title..... Gonadotropin-Releasing  
Hormone Agonists – Central Precocious  
Puberty

# Gonadotropin-Releasing Hormone Agonists – Central Precocious Puberty

- Fensolvi® (leuprolide acetate subcutaneous injection, extended-release – Tolmar)
- Lupron Depot-Ped® (leuprolide acetate depot intramuscular injection – AbbVie)
- Triptodur™ (triptorelin intramuscular injection, extended-release – Azurity)

---

### **INSTRUCTIONS FOR USE**

*The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.*

---

## OVERVIEW

Fensolvi, Lupron Depot-Ped, and Triptodur are gonadotropin-releasing hormone (GnRH) agonists indicated for the treatment of **central precocious puberty** in pediatric patients.<sup>1-3</sup>

GnRH agonists can also be used off-label for the treatment of **gender-dysphoric/gender-incongruent persons** to suppress physical changes of puberty and gonadal function.<sup>7,8</sup> Pubertal hormonal suppression should typically be initiated after the adolescent first exhibits physical changes of puberty (Tanner stages G2/B2). Long-acting GnRH analogs are the currently preferred treatment option. An advantage to using a GnRH analog is that the effects can be reversed; pubertal suppression can be discontinued if the individual no longer wishes to transition. Upon discontinuation of therapy, spontaneous pubertal development has been shown to resume. GnRH analogs can also be used in patients during late puberty to suppress the hypothalamic-pituitary-gonadal axis to allow for lower doses of cross-sex hormones.<sup>9</sup> In addition to use in adolescents, GnRH analog therapy is also used in adults, particularly male-to-female patients.<sup>10</sup>

## Dosing Information

Fensolvi is administered by a subcutaneous injection and both Lupron Depot-Ped and Triptodur are administered by intramuscular injection.<sup>1-3</sup> Fensolvi is administered once every 6 months, Lupron Depot-Ped is administered once a month, once every 3 months (12 weeks), or once every 6 months (24 weeks), and Triptodur is administered once every 24 weeks. There are no specific dosing recommendations for off-label use of Fensolvi, Lupron Depot-Ped, or Triptodur. Therefore, the FDA-approved dosing in the product labeling for approved uses has been cited for off-label uses. Treatment decisions, including duration of therapy, are individualized with careful consideration of the risks and benefit of the selected regimen.

## Guidelines

The standard of care for central precocious puberty is GnRH agonists.<sup>4-6</sup> The European Society for Paediatric Endocrinology and the Lawson Wilkins Pediatric Endocrine Society convened a consensus conference to review the use of GnRH agonists in pediatric patients with central precocious puberty (2009).<sup>4</sup> The panel noted that the available GnRH agonists (including leuprolide and triptorelin) are effective, despite different routes of administration, dosing, and duration of action. In addition, the various GnRH agonists are well-tolerated in children and adolescents. An update by an International Consortium (2019) notes the lack of prospective comparative studies to establish differences in efficacy (if any) among the various GnRH agonists.<sup>5</sup> The Consortium does not prefer one GnRH agonist over another. Discontinuation of GnRH agonist therapy should be individualized, based on the patient's readiness for resumption of puberty, recent growth rates, and shifts in height prediction.

## Coverage Policy

The use of leuprolide for oncology and infertility indications are addressed in separate coverage policies. Please refer to the related coverage policy (Oncology Medications, Infertility Injectables).

Coverage for treatment of gender dysphoria varies across plans. Coverage of drugs for hormonal therapy, as well as whether the drug is covered as a medical or a pharmacy benefit, varies across plans. Refer to the customer's benefit plan document for coverage details. In addition, coverage for treatment of gender dysphoria, including gender reassignment surgery and related services may be governed by state and/or federal mandates.

## POLICY STATEMENT

Prior Authorization is required for benefit coverage of gonadotropin-releasing hormone agonists (Fensolvi, Lupron Depot-Ped, and Triptodur). All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of gender-dysphoric/gender-incongruent persons treated with Fensolvi, Lupron Depot-Ped, or Triptodur as well as the monitoring required for adverse events and long-term efficacy, approval requires that the product be prescribed by or in consultation with a physician who specializes in this condition.

**Gonadotropin-releasing hormone agonists are considered medically necessary when ONE of the following are met (1 or 2):**

**FDA-Approved Indication**

- 1. Central Precocious Puberty.** Approve the requested gonadotropin-releasing hormone agonist for 1 year when ALL the following criteria are met:
  - A.** Onset of secondary sexual characteristics earlier than 8 years in females and 9 years in males
  - B.** Diagnosis is confirmed by ONE of the following (i or ii):
    - i. Pubertal basal level of luteinizing hormone (LH) greater than or equal to 0.2 mIU/mL
    - ii. Pubertal luteinizing hormone (LH) response to a GnRH stimulation test
  - C.** Preferred product criteria is met for the product(s) as listed in the below table(s).

**Employer Plans:**

Product	Criteria
<b>Lupron Depot-Ped</b> (leuprolide acetate depot intramuscular injection)	<b>Central Precocious Puberty.</b> Approve for 1 year when ONE of the following criteria are met (1 or 2): <ol style="list-style-type: none"> <li>1. Patient has tried ONE of Triptodur or Fensolvi</li> <li>2. Patient &lt; 2 years of age</li> </ol>

**Individual and Family Plans:**

Product	Criteria
<b>Lupron Depot-Ped</b> (leuprolide acetate depot intramuscular injection)	<b>Central Precocious Puberty.</b> Approve for 1 year when ONE of the following criteria are met (1 or 2): <ol style="list-style-type: none"> <li>1. Patient has tried ONE of Triptodur or Fensolvi</li> <li>2. Patient &lt; 2 years of age</li> </ol>

**Dosing.** Approve ONE of the following dosing regimens (A, B, or C):

- A)** Fensolvi: Approve up to one injection (45 mg) given subcutaneously once every 6 months; OR
- B)** Lupron Depot-Ped: Approve ONE of the following doses (i, ii, iii, iv, or v); OR
  - i.** 1-month depot, ≤ 25 kg: Approve up to one 1-month depot (7.5 mg) given intramuscularly (IM) once every month; OR
  - ii.** 1-month depot, > 25 to 37.5 kg: Approve up to one 1-month depot (11.25 mg) given IM once every month; OR
  - iii.** 1-month depot, > 37.5 kg: Approve up to one 1-month depot (15 mg) given IM once every month; OR
  - iv.** 3-month depot: Approve up to one 3-month depot (11.25 mg or 30 mg) given IM once every 3 months (12 weeks); OR

- v. 6-month depot: Approve up to one 6-month depot (45 mg) given IM once every 6 months (24 weeks); OR
- C) Triptodur: Approve up to one injection (22.5 mg) given IM once every 24 weeks.

**Other Uses with Supportive Evidence**

**2. Gender-Dysphoric/Gender-Incongruent Persons; Persons Undergoing Gender Reassignment (Female-To-Male or Male-To-Female).** Approve the requested gonadotropin-releasing hormone agonist for 1 year if prescribed by or in consultation with an endocrinologist or a physician who specializes in the treatment of transgender patients.

**Dosing.** Approve ONE of the following dosing regimen (A, B, or C):

- A) Fensolvi: Approve up to one injection (45 mg) given subcutaneously once every 6 months; OR
- B) Lupron Depot-Ped: Approve ONE of the following doses (i, ii, or iii); OR
  - i. 1-month depot: Approve up to one 1-month depot (7.5 mg, 11.25 mg, or 15 mg) given intramuscularly (IM) once every month; OR
  - ii. 3-month depot: Approve up to one 3-month depot (11.25 mg or 30 mg) given IM once every 3 months (12 weeks); OR
  - iii. 6-month depot: Approve up to one 6-month depot (45 mg) given IM once every 6 months (24 weeks); OR
- C) Triptodur: Approve up to one injection (22.5 mg) given IM once every 24 weeks.

**Conditions Not Covered**

**Gonadotropin-releasing hormone agonists for any other use is considered not medically necessary, including the following (this list may not be all inclusive; criteria will be updated as new published data are available):**

**1. Peripheral Precocious Puberty (Also Known as GnRH-Independent Precocious Puberty).** Children with peripheral precocious puberty do not respond to GnRH agonist therapy.<sup>4</sup> Treatment is directed at removing or blocking the production and/or response to the excess sex steroids, depending on the cause (e.g., surgically removing human chorionic gonadotropin-secreting tumors or using glucocorticoids to treat defects in adrenal steroidogenesis [such as classic congenital adrenal hyperplasia]).

Receipt of sample product does not satisfy any criteria requirements for coverage.

**Coding Information**

**Note:**

- 1) This list of codes may not be all-inclusive.
- 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

**Considered Medically Necessary when criteria in the applicable policy statements listed above are met:**

<b>HCPCS Codes</b>	<b>Description</b>
J1950	Injection, leuprolide acetate (for depot suspension), per 3.75 mg
J1951	Injection, leuprolide acetate for depot suspension (Fensolvi), 0.25 mg

HCPCS Codes	Description
J3316	Injection, triptorelin, extended release, 3.75 mg

## References

1. Lupron Depot-Ped® [prescribing information]. North Chicago, IL; AbbVie; September 2025.
2. Triptodur™ [prescribing information]. Woburn, MA: Azurity; September 2025.
3. Fensolvi® [prescribing information]. Fort Collins, CO: Tolmar; September 2025.
4. Carel JC, Eugster EA, Rogol A, et al. Consensus statement on the use of gonadotropin-releasing hormone analogs in children. *Pediatrics*. 2009;123(4):e752-62.
5. Krishna KB, Fuqua JS, Rogol AD, et al. Use of gonadotropin-releasing hormone analogs in children: update by an international consortium. *Horm Res Paediatr*. 2019;91:357-372.
6. Eugster EA. Treatment of central precocious puberty. *J Endo Soc*. 2019;3:965-972.
7. Hembree WC, Cohen-Kettenis PT, Gooren L, et al. Endocrine treatment of gender-dysphoric/gender-incongruent persons: an Endocrine Society clinical practice guidelines. *J Clin Endocrinol Metab*. 2017;102:3869-3903.
8. World Professional Association for Transgender Health (WPATH). Standards of Care for the health of transgender and gender diverse people (version 8). Available at: Standards of Care for the Health of Transgender and Gender Diverse People, Version 8 (tandfonline.com). Accessed on October 16, 2025.
9. Rosenthal SM. Approach to the patient: transgender youth: endocrine considerations. *J Clin Endocrine Metab*. 2014;99:4379-4389.
10. Spack NP. Management of transgenderism. *JAMA*. 2013;309:478-484.

## Revision Details

Type of Revision	Summary of Changes	Date
New	New policy.	2/15/2026

The policy effective date is in force until updated or retired.

---

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2026 The Cigna Group.