

Is the requested medication for a chronic or long-term condition for which the prescription medication may be necessary for the life of the patient? Yes No

Diagnosis (check all that apply to your patient):

- acquired hemophilia
- acquired inhibitor titer to Factor VIII
- acquired inhibitors to factors XI or XII
- bleeding associated with liver disease
- coagulation factor X deficiency
- congenital fibrinogen deficiency (factor I deficiency)-afibrinogenemia
- congenital fibrinogen deficiency (factor I deficiency)-hypofibrinogenemia
- congenital fibrinogen deficiency (factor I deficiency)-dysfibrinogenemia
- congenital factor VII (FVII) deficiency
- congenital factor XIII A-subunit deficiency
- congenital factor XIII B-subunit deficiency
- congenital Factor XIII deficiency
- factor II deficiency
- factor VIII deficiency (hemophilia A)
- factor IX deficiency (hemophilia B)
- factor X deficiency
- factor XIII deficiency
- Glanzmann's thrombasthenia
- hemophilia A
- hemophilia A (congenital factor VIII deficiency)
- hemophilia A with inhibitors
- hemophilia B
- hemophilia B with inhibitors
- hereditary antithrombin deficiency (antithrombin III deficiency, AT III deficiency)
- hereditary Factor X deficiency
- inhibitors to factors XI or XII
- severe von Willebrand disease (VWD)
- mild or moderate von Willebrand disease (VWD)
- All other indications or diagnoses

Clinical Information

****FEIBA, Obizur, SEVENFACT: These drugs requires supportive documentation (chart notes, lab/test results, etc) be attached with this request****

For AlphaNineSD, Alprolix, BeneFIX, Idelvion, Ixinity, Profilnine, Rebinyn and Rixubis only:

Is the requested product being prescribed by or in consultation with a hemophilia specialist? Yes No

(if BeneFIX, Ixinity, Rixubis) Is the requested dosing for up to 200 IU per kg intravenously no more frequently than once daily for immune tolerance therapy (also known as immune tolerance induction)? Yes No

(if no) Is the requested dosing for up to 100 IU per kg no more frequently than twice weekly for routine prophylaxis? Yes No

(if no) Is the requested dosing for up to 100 IU per kg intravenously no more frequently than once every 12 hours for up to 10 days per episode for on-demand treatment of and control of bleeding episodes? Yes No

(if no) Is the requested dosing for up to 100 IU per kg intravenously no more frequently than once every 8 hours for up to 10 days per procedure for perioperative management? Yes No

(if Alprolix, Idelvion, Rebinyn) Is the requested dosing for up to 100 IU per kg intravenously at an interval no more frequently than once weekly for routine prophylaxis? Yes No

(if no) Is the requested dosing for up to 100 IU per kg intravenously no more frequently than once every 6 hours for up to 10 days per episode for on-demand treatment of and control of bleeding episodes? Yes No

(if no) Is the requested dosing for up to 100 IU per kg intravenously no more frequently than once every 24 hours for up to 10 days per procedure for perioperative management? Yes No

(if AlphaNine SD, Profilnine) Is the requested dosing for up to 200 IU per kg intravenously no more frequently than once daily for immune tolerance therapy (also known as immune tolerance induction)? Yes No

(if no) Is the requested dosing for up to 50 IU per kg no more frequently than twice weekly for routine prophylaxis? Yes No

(if no) Is the requested dosing for up to 100 IU per kg intravenously no more frequently than twice daily for up to 10 days for on-demand treatment of and control of bleeding episodes and perioperative management? Yes No

For Advate, Afstyla, Kogenate FS, Kovaltry, Novoeight, Nuwiq, Recombinate, or Xyntha:

Is the requested product being prescribed by or in consultation with a hemophilia specialist? Yes No

Is the requested dosing for up to 60 IU per kg intravenously no more frequently than every other day (three or four times weekly) for routine prophylaxis? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing up to 50 IU per kg intravenously no more frequently than once every 6 hours for up to 10 days per episode for on-demand treatment and control of bleeding episodes? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing for up to 60 IU per kg intravenously no more frequently than once every 6 hours for up to 10 days per procedure for perioperative management? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing for up to 200 IU per kg intravenously no more frequently than once daily for immune tolerance therapy (also known as immune tolerance induction)? Please be sure to provide the patient's weight in kg. Yes No

For Adynovate, Elocate, Esperoct, or Jivi:

Is the requested product being prescribed by or in consultation with a hemophilia specialist? Yes No

Is the requested dosing for up to 100 IU per kg intravenously no more frequently than twice weekly for routine prophylaxis? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing up to 65 IU per kg intravenously no more frequently than once every 8 hours for up to 10 days per episode for on-demand treatment and control of bleeding episodes? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing for up to 65 IU per kg intravenously no more frequently than once every 6 hours for up to 10 days per procedure for perioperative management? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing for up to 200 IU per kg intravenously no more frequently than once daily for immune tolerance therapy (also known as immune tolerance induction)? Please be sure to provide the patient's weight in kg. Yes No

For Alphanate, Humate-P, or Wilate only:

Is the requested product being prescribed by or in consultation with a hemophilia specialist? Yes No

(if hemophilia A) Is the requested dosing for up to 50 IU per kg intravenously no more frequently than every other day (three or four times weekly) for routine prophylaxis? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing up to 50 IU per kg intravenously no more frequently than once every 6 hours for up to 10 days per episode or procedure for on-demand treatment and control of bleeding episodes or perioperative management? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing for up to 200 IU per kg intravenously no more frequently than once daily for immune tolerance therapy (also known as immune tolerance induction)? Please be sure to provide the patient's weight in kg. Yes No

(if vWD) What type of von Willebrand Disease (vWD) does the patient have?

- Type 1
- Type 2
- Type 3

(if Type 1) Is the requested medication being used for the treatment of bleeding episodes? Yes No

(if yes) Has the patient tried desmopressin injection (DDAVP injection)? Yes No

(if no) Does the patient have a severe bleeding phenotype? Yes No

(if no) Does the patient have very low vWF (von Willebrand Factor) levels? Yes No

(if no OR not for treatment of bleeding episodes) Does the patient have a history of inadequate response to injectable desmopressin in the past or has safety concerns with use of desmopressin injection (DDAVP injection)? Yes No

(if vWD) Is the requested dosing for up to 80 IU VWF:Rco (Von Willebrand Factor activity as measured with the Ristocetin cofactor assay) per kg intravenously no more frequently than once every 8 hours for up to 10 days per episode or procedure for on-demand treatment and control of bleeding episodes or perioperative management? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing for up to 40 IU VWF:Rco (Von Willebrand Factor activity as measured with the Ristocetin cofactor assay) per kg intravenously no more frequently than once every 2 days for routine prophylaxis? Please be sure to provide the patient's weight in kg. Yes No

(if ATryn) Is ATryn being used for the prevention of perioperative or peripartum events? Yes No

(if Fibryga or RiaSTAP) Has the patient had testing showing prolonged activated partial thromboplastin time and prothrombin time at baseline, as defined by the laboratory reference values? Yes No

(if Fibryga or RiaSTAP) Has the patient had testing showing lower than normal plasma functional and antigenic fibrinogen levels at baseline, as defined by the laboratory reference values? Yes No

(if Fibryga or RiaSTAP) Is this medication being prescribed by, or in consultation with, a hematologist? Yes No

(if Fibryga or RiaSTAP) Will both Fibryga and RiaSTAP be taken together at the same time? Yes No

(If yes) Please provide the clinical rationale for concurrent use:

(if Coagadex)

For which of the following is this drug being used?

- Peri-operative management of bleeding in individuals with mild, moderate, or severe hereditary Factor X deficiency
- Routine prophylaxis to reduce the frequency of bleeding episodes
- Treatment of bleeding episodes
- Other

(if other) Please provide clinical rationale for the use of this drug in your patient.

For Altuviio only:

Is Altuviio being used in at least one of the following scenarios: 1) Routine prophylaxis; or 2) On-demand treatment and control of bleeding episodes; or 3) Perioperative management of bleeding?

- Routine prophylaxis
- On-demand treatment and control of bleeding episodes
- Perioperative management of bleeding
- None of the above scenarios apply

Is the medication being prescribed by or in consultation with a hemophilia specialist? Yes No

Is the patient currently receiving Altuviio or has received Altuviio in the past? Yes No

(if no) Has the patient received Factor VIII therapy in the past? Yes No

(if yes) Has Factor VIII inhibitor testing been performed within the past 30 days? Yes No

(if yes) Does the patient have a positive test for Factor VIII inhibitors greater than or equal to 1.0 Bethesda units/mL? Yes No

(if currently receiving or has received Altuviio in the past) Has Factor VIII inhibitor testing been performed within the past 365 days? Yes No

(if yes) Does the patient have a positive test for Factor VIII inhibitors greater than or equal to 1.0 Bethesda units/mL? Yes No

(if no Factor VIII inhibitor testing in past 365 days OR no positive test for Factor VIII inhibitors 1.0 Bethesda units/ml or more) According to the prescriber, does the patient have clinical manifestations suggesting the presence of Factor VIII inhibitors? Please Note: Inhibitors may be present if bleeding is not well controlled, there is decreased responsiveness to Factor VIII therapy, and/or if expected Factor VIII activity plasma levels are not achieved. Yes No

If Hemofil M or Koate:

Is the requested product being prescribed by or in consultation with a hemophilia specialist? Yes No

Is the requested dosing for up to 50 IU per kg intravenously no more frequently than every other day (three or four times weekly) for routine prophylaxis? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing up to 50 IU per kg intravenously no more frequently than once every 6 hours for up to 10 days per episode or procedure for on-demand treatment and control of bleeding episodes or perioperative management? Please be sure to provide the patient's weight in kg. Yes No

(if no) Is the requested dosing for up to 200 IU per kg intravenously no more frequently than once daily for immune tolerance therapy (also known as immune tolerance induction)? Please be sure to provide the patient's weight in kg. Yes No

(if Vonvendi)

For which of the following is this drug being used?

- On demand treatment and control of bleeding episodes
- Peri-operative management
- Routine prophylaxis
- None of the above

(if Coagadex, Corifact, Tretten, Vonvendi) Is the requested medication being prescribed by, or in consultation with, a hematologist? Yes No

(if Hemlibra) Is there documentation that your patient has one of the following?

- factor XIII inhibitors
- mild or moderate hemophilia (defined as factor VIII level of 1% to less than 40%)
- severe hemophilia defined as pre-treatment factor VIII level less than 1%
- none of the above

(if mild/moderate) Which of the following applies to your patient? Please provide documentation

- 1 or more episodes of bleeding into the central nervous system or other serious, life-threatening bleed
- 1 or more episodes of bleeding into large joint (ankles, knees, hips, elbows, shoulders) and age 3 years or younger
- 2 or more episodes of bleeding into large joints (ankles, knees, hips, elbows, shoulders)
- presence of joint disease documented by physical examination and plain radiographs of the affected joints
- none of the above

(if Hemlibra) Is Hemlibra being used for routine prophylaxis to prevent or reduce the frequency of bleeding episodes? Yes No

(if no) Please specify the use for which Hemlibra is being prescribed.

(if Obizur) For which of the following is the drug requested being used?

- treatment of current active bleed
- prevention of excessive bleeding during and/or following surgery
- routine prophylaxis
- as needed dosing for future bleeds
- other

(if surgery) What is the date of surgery?

(if as needed dosing) What is the approximate number of bleeds requiring factor treatment per month?

(if other) Please provide clinical rationale for the use of this drug in your patient.

(if Obizur) (if acquired hemophilia) Has there been documentation provided of autoimmune inhibitory antibodies to human factor VIII? Yes No

(if Obizur) Does the patient have a diagnosis of either congenital hemophilia A or von Willebrand's disease? Yes No

(if NovoSeven RT) (if Glanzmann's thrombasthenia) Is the patient refractory to platelet transfusions? Yes No

NovoSeven RT or Sevenfact:

(if NovoSeven RT and Glanzmann's thrombasthenia) Is the requested medication being prescribed by, or in consultation with, a hematologist? Yes No

(if Sevenfact) Is the requested medication being prescribed by or in consultation with a hemophilia specialist? Yes No

(if Hemophilia A with inhibitors) (If Hemophilia A with inhibitors) Is documentation being provided that the patient has a positive inhibitor titer greater than or equal to 5 Bethesda Units? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

(if no) Is documentation being provided that the patient has a history of an inhibitor with anamnestic response to Factor VIII replacement therapy, which, according to the prescriber, precludes the use of Factor VIII replacement to treat bleeding episodes? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

(if no) Is documentation being provided that the patient has a history of an inhibitor with refractory hemostatic response to increased Factor VIII dosing, which, according to the prescriber, precludes the use of Factor VIII replacement to treat bleeding episodes? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

((if Hemophilia B with inhibitors) Is documentation being provided that the patient has a positive inhibitor titer greater than or equal to 5 Bethesda Units? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

(if no) Is documentation being provided that the patient has a history of an inhibitor with anamnestic response to Factor IX replacement therapy, which, according to the prescriber, precludes the use of Factor IX replacement to treat bleeding episodes? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

(if no) Is documentation being provided that the patient has a history of an inhibitor with refractory hemostatic response to increased Factor IX dosing, which, according to the prescriber, precludes the use of Factor IX replacement to treat bleeding episodes? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

(if NovoSeven RT and hemophilia [acquired, A/B with inhibitors]) Is the requested medication being prescribed by or in consultation with a hemophilia specialist? Yes No

(if Thrombate III) Is Thrombate III being used to treat or prevent pulmonary or deep vein embolisms (PE, DVT)? Yes No

If yes, please include the most recent clinical notes.

(if Thrombate III) Is your patient undergoing a surgical or obstetrical procedure? Yes No

For FEIBA only:

(if Hemophilia A with inhibitors):

Is documentation being provided that the patient has a positive inhibitor titer greater than or equal to 5 Bethesda Units? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

Is documentation being provided that the patient has a history of an inhibitor with anamnestic response to Factor VIII replacement therapy, which, according to the prescriber, precludes the use of Factor VIII replacement to treat bleeding episodes? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

Is documentation being provided that the patient has a history of an inhibitor with refractory hemostatic response to increased Factor VIII dosing, which, according to the prescriber, precludes the use of Factor VIII replacement to treat bleeding episodes? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

(if Hemophilia B with inhibitors):

Is documentation being provided that the patient has a positive inhibitor titer greater than or equal to 5 Bethesda Units? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

Is documentation being provided that the patient has a history of an inhibitor with anamnestic response to Factor IX replacement therapy, which, according to the prescriber, precludes the use of Factor IX replacement to treat bleeding episodes? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

Is documentation being provided that the patient has a history of an inhibitor with refractory hemostatic response to increased Factor IX dosing, which, according to the prescriber, precludes the use of Factor IX replacement to treat bleeding episodes? PLEASE NOTE: Medical documentation specific to your response must be attached to this case or your request may be denied. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. Yes No

(if Hemophilia A OR Hemophilia B with Inhibitors):

Is the requested medication being prescribed by or in consultation with a hemophilia specialist? Yes No

Additional pertinent information:

Attestation: I attest the information provided is true and accurate to the best of my knowledge. I understand that the Health Plan or insurer its designees may perform a routine audit and request the medical information necessary to verify the accuracy of the information reported on this form.

Prescriber Signature: _____ **Date:** _____

Save Time! Submit Online at: www.covermymeds.com/main/prior-authorization-forms/cigna/ or via SureScripts in your EHR.

Our standard response time for prescription drug coverage requests is 5 business days. If your request is urgent, it is important that you call us to expedite the request. View our Prescription Drug List and Coverage Policies online at cigna.com.

V050126

"Cigna" is a registered service mark, and the "Tree of Life" logo is a service mark, of Cigna Intellectual Property, Inc., licensed for use by Cigna Corporation and its operating subsidiaries. All products and services are provided by or through such operating subsidiaries and not by Cigna Corporation. Such operating subsidiaries include, for example, Cigna Health and Life Insurance Company and Cigna Health Management, Inc. Address: Cigna Pharmacy Services, PO Box 42005, Phoenix AZ 85080-2005