



Xolair (omalizumab)

Fax completed form to: (855) 840-1678
 If this is an URGENT request, please call (800) 882-4462
 (800.88.CIGNA)

PHYSICIAN INFORMATION			PATIENT INFORMATION		
* Physician Name:			*Due to privacy regulations we will not be able to respond via fax with the outcome of our review unless all asterisked (*) items on this form are completed.*		
Specialty:	* DEA, NPI or TIN:				
Office Contact Person:			* Patient Name:		
Office Phone:			* Cigna ID:		* Date of Birth:
Office Fax:			* Patient Street Address:		
Office Street Address:			City:		State:
City:			State:		Zip:
State:			Patient Phone:		
Zip:					
Urgency: <input type="checkbox"/> Standard <input type="checkbox"/> Urgent (In checking this box, I attest to the fact that applying the standard review time frame may seriously jeopardize the customer's life, health, or ability to regain maximum function)					
Medication requested: <input type="checkbox"/> Xolair 150mg vial <input type="checkbox"/> Xolair 75mg/0.5ml syringe or autoinjector <input type="checkbox"/> Xolair 150mg/ml syringe or autoinjector <input type="checkbox"/> Xolair 300mg/2ml syringe or autoinjector <input type="checkbox"/> Other (<i>please specify</i>):					
Directions for use, dose, and quantity:			Duration of therapy:		
Frequency of therapy:					
J-Code:			ICD10:		
Where will this medication be obtained? <input type="checkbox"/> Accredo Specialty Pharmacy** <input type="checkbox"/> Hospital Outpatient <input type="checkbox"/> Retail pharmacy <input type="checkbox"/> Other (<i>please specify</i>): <input type="checkbox"/> Home Health / Home Infusion vendor <input type="checkbox"/> Physician's office stock (billing on a medical claim form) <i>**Cigna's nationally preferred specialty pharmacy</i> 					
<i>**Medication orders can be placed with Accredo via E-prescribe - Accredo (1620 Century Center Pkwy, Memphis, TN 38134-8822 NCPDP 4436920), Fax 888.302.1028, or Verbal 866.759.1557</i>					
Facility and/or doctor dispensing and administering medication: Facility Name: _____ State: _____ Tax ID#: _____ Address (City, State, Zip Code): _____					
Where will this drug be administered? <input type="checkbox"/> Patient's Home <input type="checkbox"/> Hospital Outpatient <input type="checkbox"/> Physician's Office <input type="checkbox"/> Other (<i>please specify</i>): 					
NOTE: Per some Cigna plans, infusion of medication MUST occur in the least intensive, medically appropriate setting. Is this patient a candidate for re-direction to an alternate setting (such as alternate infusion site, physician's office, home) with assistance of a Specialty Care Options Case Manager? <input type="checkbox"/> Yes <input type="checkbox"/> No (<i>provide medical necessity rationale</i>):					
Is the requested medication for a chronic or long-term condition for which the prescription medication may be necessary for the life of the patient? <input type="checkbox"/> Yes <input type="checkbox"/> No 					

Clinical Data:

What diagnosis is Xolair being used to treat?

- atopic dermatitis
- asthma
- Chronic Spontaneous Urticaria
- Chronic Rhinosinusitis with Nasal Polyps (CRSwNP)
- eosinophilic gastroenteritis (EG), eosinophilic esophagitis (EE), or eosinophilic colitis
- Immunoglobulin (Ig)E-Mediated Food Allergy
- Treatment of latex allergy in healthcare workers with occupational latex allergy
- All other indications of diagnoses

Will Xolair be used in combination with another Monoclonal Antibody Therapy? Monoclonal antibody therapies are Adbry (tralokinumab-ldrm subcutaneous injection), Cinqair (reslizumab intravenous infusion), Dupixent (dupilumab subcutaneous injection), Ebglyss (lebrikizumab-lbkz SC injection), Fasentra (benralizumab subcutaneous injection), Nemluvio (nemolizumab-ilto SC injection), Nucala (mepolizumab subcutaneous injection), or Tezspire (tezepelumab-ekko subcutaneous injection). Yes No

(if asthma, CRSwNP, or IgE Mediated Food Allergy) Does the patient have a baseline immunoglobulin E (IgE) level greater than or equal to 30 IU/mL? Notes: Please Note: "Baseline" is defined as prior to receiving any treatment with Xolair or another monoclonal antibody therapy that may lower IgE levels (for example, Dupixent [dupilumab subcutaneous injection], Tezspire [tezepelumab-ekko subcutaneous injection]). Yes No

(if asthma) Is the requested medication being prescribed by or in consultation with an allergist, immunologist, or pulmonologist? Yes No

(if Chronic Spontaneous Urticaria) Is the requested medication being prescribed by, or in consultation with, an allergist, immunologist, or dermatologist? Yes No

(if CRSwNP) Is the requested medication being prescribed by, or in consultation with, an allergist, immunologist, or otolaryngologist (ear, nose, and throat [ENT] physician specialist)? Yes No

(if IgE Mediated Food Allergy) Is the requested medication prescribed by, or in consultation with, an allergist or an immunologist? Yes No

Is the patient currently receiving Xolair? Yes No

If currently receiving:

(if asthma, currently receiving) Has the patient already received at least 4 months of therapy with Xolair? Please Note: Answer No if the patient has received less than 4 months of therapy or if the patient is restarting therapy with Xolair. Yes No

(if Chronic Spontaneous Urticaria, CRSwNP currently receiving) Has the patient already received at least 6 months of therapy with Xolair? Please Note: Answer No if the patient has received less than 6 months of therapy or if the patient is restarting therapy with Xolair. Yes No

(if asthma, currently receiving) Will the patient continue to receive therapy with either one inhaled corticosteroid or one inhaled corticosteroid-containing combination inhaler? Yes No

((if asthma, currently receiving) Has the prescriber determined that the patient has responded to therapy? Please Note: Examples of a response to Xolair therapy are decreased asthma exacerbations; decreased asthma symptoms; decreased hospitalizations, emergency department, urgent care, or medical clinic visits due to asthma; decreased reliever/rescue medication use; and improved lung function parameters. Yes No

(if Chronic Spontaneous Urticaria, CRSwNP currently receiving) Has the patient already received at least 6 months of therapy with Xolair? Please Note: Answer No if the patient has received less than 6 months of therapy or if the patient is restarting therapy with Xolair. Yes No

(if Chronic Spontaneous Urticaria, currently receiving) Has the patient experienced a beneficial clinical response, defined by one of the following: a. Decreased itch severity; b. Decreased number of hives; or c. Decreased size of hives? Yes No

(if CRSwNP, currently receiving) Will the patient continue to receive therapy with an intranasal corticosteroid? Yes No

(if CRSwNP, currently receiving) Has the patient responded to therapy as determined by the prescriber (Examples of a response include reduced nasal polyp size, improved nasal congestion, reduced sinus opacification, decreased sino-nasal symptoms, and/or improved sense of smell)? Yes No

If asthma and initial therapy, restarting therapy, or currently receiving less than 4 months of therapy:

Did/Does the patient have a forced expiratory volume in 1 second (FEV1) of less than 80% predicted? (Note: The reduced FEV1 should not be due to smoking-related chronic obstructive pulmonary disease. Also the above lung function criteria may be met at anytime prior to or during asthma treatment.) Yes No

(if yes) Did/Does the patient have an FEV1/forced vital capacity (FVC) of less than 0.80? (Note: The above lung function criteria may be met at anytime prior to or during asthma treatment.) Yes No

(if no, and age 6-11 years) Did/Does the patient have an increase of greater than or equal to 12% in FEV1 following the administration of a standard dose of a short-acting bronchodilator? (Note: The above lung function criteria may be met at anytime prior to or during asthma treatment.) Yes No

(if no, and age 6-11 years) Did/Does the patient have an increase of greater than or equal to 12% in FEV1 between prescriber visits? (Note: The above lung function criteria may be met at anytime prior to or during asthma treatment.) Yes No

(if no, and age 6-11 years) Did/Does the patient have an increase of greater than or equal to 12% in FEV1 from baseline to after at least 4 weeks of asthma treatment? (Note: The above lung function criteria may be met at anytime prior to or during asthma treatment.) Yes No

(if no FEV1/FVC less than 0.80, and age 12 years and older) Did/Does the patient have an increase of greater than or equal to 12% AND greater than or equal to 200 mL in FEV1 following the administration of a standard dose of a short-acting bronchodilator? (Note: The above lung function criteria may be met at anytime prior to or during asthma treatment.) Yes No

(if no, and age 12 years and older) Did/Does the patient have an increase of greater than or equal to 12% AND greater than or equal to 200 mL in FEV1 between prescriber visits? (Note: The above lung function criteria may be met at anytime prior to or during asthma treatment.) Yes No

(if no, and age 12 years and older) Did/Does the patient have an increase of greater than or equal to 12% AND greater than or equal to 200 mL in FEV1 from baseline to after at least 4 weeks of asthma treatment? (Note: The above lung function criteria may be met at anytime prior to or during asthma treatment.) Yes No

(if no FEV1/FVC less than 0.8, AND no increase greater than or equal to 12% in FEV1 between prescriber visits/from baseline to after at least 4 weeks of asthma treatment) Did/Does the patient have a positive exercise challenge test? Note: The above lung function criteria may be met at anytime prior to or during asthma treatment. Yes No

(if no) Did/Does the patient have a positive bronchial challenge test? Note: The above lung function criteria may be met at anytime prior to or during asthma treatment. Yes No

Has the patient had a baseline positive skin test or in vitro test (that is, a blood test) for allergen-specific immunoglobulin E (IgE) for one or more perennial aeroallergens and/or for one or more seasonal aeroallergens? Notes: Please Note: "Baseline" is defined as prior to receiving any Xolair or another monoclonal antibody therapy that may interfere with allergen testing (for example, Dupixent and Tezspire). Examples of perennial aeroallergens are house dust mite, animal dander, cockroach, feathers, and mold spores. Examples of seasonal aeroallergens are grass, pollen, and weeds Yes No

Has the patient received at least 3 consecutive months of combination therapy with BOTH of the following: an inhaled medium- or high-dose corticosteroid AND at least one additional asthma controller or asthma maintenance medication. Notes: Please Note: Examples of additional asthma controller or asthma maintenance medications are inhaled long-acting beta2-agonists, inhaled long-acting muscarinic antagonists, and monoclonal antibody therapies for asthma (for example, Xolair, Cinqair [reslizumab intravenous infusion], Dupixent, Fasenra [benralizumab subcutaneous injection], Nucala [mepolizumab subcutaneous injection], and Tezspire). Please Note: Use of a combination inhaler containing both a medium- or high- dose inhaled corticosteroid and additional asthma controller or asthma maintenance medication(s) would fulfil the requirement. Yes No

Does the patient have asthma that is uncontrolled or was uncontrolled at baseline as defined by the patient experiencing two or more asthma exacerbations requiring treatment with systemic corticosteroids in the previous year? Notes: Please Note: "Baseline" is defined as prior to receiving Xolair or another monoclonal antibody therapy for asthma. Examples of monoclonal antibody therapies for asthma include Cinqair, Dupixent, Fasenra, Nucala, Tezspire, and Xolair. Yes No

(if no) Does the patient have asthma that is uncontrolled or was uncontrolled at baseline as defined by the patient experiencing one or more asthma exacerbation(s) requiring a hospitalization, an emergency department visit, or an urgent care visit in the previous year? Notes: Please Note: "Baseline" is defined as prior to receiving Xolair or another monoclonal antibody therapy for asthma. Examples of monoclonal antibody therapies for asthma include Cinqair, Dupixent, Fasenra, Nucala, Tezspire, and Xolair. Yes No

(if no) Does the patient have asthma that is uncontrolled or was uncontrolled at baseline as defined by the patient having asthma that worsens upon tapering of oral corticosteroid therapy? Notes: Please Note: "Baseline" is defined as prior to receiving Xolair or another monoclonal antibody therapy for asthma. Examples of monoclonal antibody therapies for asthma include Cinqair, Dupixent, Fasenra, Nucala, Tezspire, and Xolair. Yes No

If Chronic Rhinosinusitis with Nasal Polyps and Initial Therapy, Restarting Therapy, or Currently Receiving for less than 6 months:

Does the patient have chronic rhinosinusitis with nasal polyps as evidenced by direct examination, endoscopy, or sinus computed tomography (CT) scan? Yes No

Has the patient had the diagnosis of chronic rhinosinusitis with nasal polyps for at least 6 months? Yes No

Has the patient experienced TWO or more of the following symptoms for at least 8 weeks: nasal congestion, nasal obstruction, nasal discharge, and/or reduction/loss of smell? Yes No

Has the patient received an intranasal corticosteroid for at least 4 weeks? Yes No

(if yes) Will the patient continue to receive therapy with an intranasal corticosteroid concomitantly with Xolair? Yes No

Has the patient had prior surgery for nasal polyps? Yes No

(if no) Has the patient received at least one course of treatment with a systemic corticosteroid within the previous year?
Please note: One course of a systemic corticosteroid is greater than or equal to 3 consecutive days of treatment or one long-acting injectable dose. Yes No

(if no) Does the patient have a contraindication to systemic corticosteroid therapy? Yes No

If Chronic Spontaneous Urticaria and Initial Therapy, Restarting Therapy, or Currently Receiving for less than 6 months:

Has the patient had urticaria for greater than or equal to 6 weeks (prior to treatment with Xolair)? Yes No

According to the prescriber, has the patient tried high-dose oral second-generation H1 antihistamine therapy? Please Note: High-dose oral second-generation H1 antihistamine therapy is the highest dose tolerated by the patient and can be up to four times the FDA-approved dose. Examples of second-generation H1 antihistamines are cetirizine, desloratadine, fexofenadine, levocetirizine, and loratadine. Yes No

If Immunoglobulin (Ig)E-Mediated Food Allergy:

Has the patient had a positive skin prick test (SPT) response to one or more foods? Yes No

(if no) Has the patient had a positive in vitro test (that is, a blood test) for IgE to one or more foods? Yes No

According to the prescriber, has the patient demonstrated signs and symptoms of a significant systemic allergic reaction to a food? (Note: Signs and symptoms of a significant systemic allergic reaction include hives, swelling, wheezing, hypotension, and gastrointestinal symptoms). Yes No

(if yes) According to the prescriber, did this reaction occur within a short period of time following a known ingestion of the food? Yes No

(if yes) Has the prescriber deemed this reaction significant enough to require a prescription for an epinephrine self-administered injectable or nasal product (Examples include EpiPen, EpiPen Jr., Auvi-Q, and generic epinephrine auto-injectors, and Neffy)? Yes No

Has the patient been prescribed an epinephrine self-administered injectable or nasal product (Examples include EpiPen, EpiPen Jr., Auvi-Q, and generic epinephrine auto-injectors, and Neffy)? Yes No

According to the prescriber, will Xolair be used in conjunction with a food allergen-avoidant diet? Yes No

Additional Pertinent Information:

Attestation: I attest the information provided is true and accurate to the best of my knowledge. I understand that the Health Plan or insurer its designees may perform a routine audit and request the medical information necessary to verify the accuracy of the information reported on this form.

Prescriber Signature: _____ **Date:** _____

Save Time! Submit Online at: www.covermymeds.com/main/prior-authorization-forms/cigna/ or via SureScripts in your EHR.

Our standard response time for prescription drug coverage requests is 5 business days. If your request is urgent, it is important that you call us to expedite the request. View our Prescription Drug List and Coverage Policies online at cigna.com.

v060126